

Please write clearly in block capitals.

Centre number

--	--	--	--	--

Candidate number

--	--	--	--

Surname

Forename(s)

Candidate signature

INTERNATIONAL A-LEVEL CHEMISTRY (9620)

Unit 5 Practical and synoptic

Monday 18 June 2018

07:00 GMT

Time allowed: 1 hour 25 minutes

Materials

For this paper you must have:

- the Periodic Table/Data Sheet, provided as an insert
- a ruler with millimetre measurements
- a scientific calculator, which you are expected to use where appropriate.

Instructions

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer the questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- All working must be shown.
- Do all rough work in this book. Cross through any work you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 60.

For Examiner's Use	
Question	Mark
1	
2	
3	
4–33	
TOTAL	



Section AAnswer **all** questions in the spaces provided.**0 1 . 1**

A student has samples of three organic compounds. The samples are not labelled but are known to be propanoic acid, butanal and pentan-1-ol.

Describe **three** test-tube reactions that can be used to identify the three organic compounds. Use a **different** test-tube reaction for each compound.

Your answer should include the reagent(s) used and the expected observation for each test.

[6 marks]

Propanoic acid _____

Butanal _____

Pentan-1-ol _____



0	1	2
---	---	---

The student mixed together samples of pentan-1-ol and propanoic acid and added a few drops of concentrated sulfuric acid.

The mixture was left in a warm place for a reaction to occur.

Write an equation for the reaction that occurs.

State an observation that would be made at the end of the reaction.

Give the IUPAC name of the organic product.

[3 marks]

Equation

Observation _____

IUPAC name _____

9


Turn over for the next question

Turn over ►



0 2

An alcohol, ROH, reacts with hydrochloric acid according to the equation



The kinetics of this reaction were investigated using this method.

- An aqueous mixture was prepared containing equal volumes of alcohol and hydrochloric acid.
- The concentration of HCl in the mixture, [HCl], was monitored by removing 5.00 cm³ samples at known times. The samples were diluted and titrated with aqueous sodium hydroxide.
- A graph was plotted of [HCl] against time.

0 2 . 1

A 5.00 cm³ sample was removed from the reaction mixture. This sample was then diluted to a volume of 25.0 cm³ using deionised water.

This diluted sample needed 11.50 cm³ of 0.255 mol dm⁻³ aqueous sodium hydroxide for complete neutralisation.

Calculate [HCl] in the sample removed from the reaction mixture.
Give your answer to the appropriate number of significant figures.


[3 marks]

[HCl] _____ mol dm⁻³



0 2 . 2 Figure 1 shows how $[\text{HCl}]$ varies with time during the reaction.

Figure 1



Determine the rate of reaction at $t = 4.0$ minutes by drawing a tangent to the curve in **Figure 1**.

[3 marks]

rate of reaction _____ $\text{mol dm}^{-3} \text{ min}^{-1}$

Turn over ►



0 2 . 3 A graph of rate of reaction against [HCl] was plotted. The graph was a straight line through the origin.

The reaction between ROH and HCl is first order with respect to ROH

Deduce the rate equation for the reaction between ROH and HCl

[1 mark]

0 2 . 4 In one calculation of [HCl] the volume of the sample removed was recorded as 5.00 cm³ but 10.00 cm³ were incorrectly removed.

Deduce the effect, if any, of this mistake on the calculated value of [HCl]

[1 mark]

0 2 . 5 The kinetics of the reverse reaction ($\text{RCl} + \text{H}_2\text{O} \rightarrow \text{ROH} + \text{HCl}$) were also investigated.
The reaction was found to be first order with respect to RCl and zero order with respect to H₂O

In one experiment, the results obtained were

$$\begin{aligned}[\text{RCl}] &= 0.107 \text{ mol dm}^{-3} \\ \text{rate} &= 0.562 \text{ mol dm}^{-3} \text{ min}^{-1}\end{aligned}$$

Calculate the value of the rate constant, k , for this reverse reaction.
Deduce its units.

[2 marks]



k _____

units _____



0 2 . 6

Hydrolysis reactions such as that between RCl and H₂O occur by different mechanisms that can be represented as shown.



Deduce which mechanism (**1** or **2**) is supported by the information given in Question **02.5**

Explain your choice.

[2 marks]

12

Turn over for the next question

Turn over ►




0 3

This question is about electrochemical cells.

Figure 2 shows a diagram of an electrochemical cell.

Figure 2



0 3 . 1

Explain how the salt bridge provides an electrical connection between the two half-cells.

[1 mark]

0 3 . 2

Suggest why potassium chloride would **not** be a suitable salt for the salt bridge in this cell.

[1 mark]

0 3 . 3



Suggest why the current in the external circuit of this cell falls to zero after the cell has operated for some time.

[1 mark]



0 3 4

A different electrochemical cell was made from a copper half-cell and a manganese half-cell.



Give the conventional representation for this cell.

Calculate the EMF of this cell.

[2 marks]

Cell representation _____

EMF _____ V

Question 3 continues on the next page

Turn over ►

Table 1 shows some standard electrode potential data.

Table 1

Electrode half-equation	E^\ominus / V
$\text{Fe}^{3+} + \text{e}^- \rightarrow \text{Fe}^{2+}$	+0.77
$\text{Cu}^{2+} + 2\text{e}^- \rightarrow \text{Cu}$	+0.34
$\text{Fe}^{2+} + 2\text{e}^- \rightarrow \text{Fe}$	-0.44
$\text{Zn}^{2+} + 2\text{e}^- \rightarrow \text{Zn}$	-0.76

Use these data to answer the following questions.

0 3 . 5

Iron filings are added to a solution that contains the ions Cu^{2+} , Fe^{3+} and Zn^{2+} , all with concentration 1.00 mol dm^{-3}

Give the formula of the metal ion in the solution that does **not** react when iron filings are added.

Give a reason for your answer.

[2 marks]

0 3 . 6

Deduce the equation for each of the two reactions that occur.

[2 marks]



Section B

Do not write
outside the
boxEach Question is followed by four responses, **A, B, C** and **D**.

For each question select the best response.

Only **one** answer per question is allowed.

For each answer completely fill in the circle alongside the appropriate answer.

CORRECT METHOD



WRONG METHODS



If you want to change your answer you must cross out your original answer as shown.

If you wish to return to an answer previously crossed out, ring the answer you now wish to select as shown.

You may do your working in the blank space around each question but this will not be marked.

0 4 Which element is in the p block of the Periodic Table?**[1 mark]****A** Technetium**B** Terbium**C** Thallium**D** Thorium**0 5** Which statement about sulfur dioxide is correct?**[1 mark]****A** It is an ionic compound.**B** It is insoluble in water.**C** It reacts with calcium hydroxide solution.**D** It reacts with hydrochloric acid.

Turn over ►



0 6 Which molecule contains the smallest bond angle?

[1 mark]


A H₂S

B BH₃


C CF₄

D PCl₅

0 7 The structures of two aromatic compounds are shown.



2-chloromethylbenzene



phenylchloromethane

2-Chloromethylbenzene is less reactive towards nucleophiles than phenylchloromethane.

This is because in 2-chloromethylbenzene

[1 mark]

A the C – Cl bond is less polar.

B the ring of delocalised electrons would be broken.

C the chlorine atom withdraws electrons from the benzene ring.

D the C – Cl bond is stronger.



0 8 Which of the following is **not** able to act as a ligand?

[1 mark]

A Cl^-

B $\text{CH}_3\text{CH}_2\text{CH}_3$

C H_2O

D $\text{H}_2\text{NCH}_2\text{CH}_2\text{NH}_2$

0 9 Which equation represents the reaction for which the energy change is the standard enthalpy of formation of sodium hydroxide?

[1 mark]

A $\text{Na(s)} + \text{H}_2\text{O(l)} \rightarrow \text{NaOH(s)} + \frac{1}{2}\text{H}_2\text{(g)}$

B $\text{Na(s)} + \frac{1}{2}\text{O}_2\text{(g)} + \frac{1}{2}\text{H}_2\text{(g)} \rightarrow \text{NaOH(s)}$

C $\text{Na(s)} + \frac{1}{2}\text{O}_2\text{(g)} + \frac{1}{2}\text{H}_2\text{(g)} \rightarrow \text{NaOH(aq)}$


D $2\text{Na(s)} + \text{H}_2\text{(g)} + \text{O}_2\text{(g)} \rightarrow 2\text{NaOH(s)}$

Turn over for the next question

Turn over ►



1 0 The diagram shows the energy profiles for a reaction, with and without using a catalyst.



Which letter represents the activation energy for the catalysed reaction?

[1 mark]

- A W
- B X
- C Y
- D Z


1 1 What is the expression for the dissociation constant (K_a) for the weak acid HX?

[1 mark]

- A $\frac{[H^+] + [X^-]}{[HX]}$
- B $\frac{[HX]}{[H^+] + [X^-]}$
- C $\frac{[HX]}{[H^+][X^-]}$
- D $\frac{[H^+][X^-]}{[HX]}$



1 2 The equation for a displacement reaction is



Which statement explains why bromine displaces iodine from potassium iodide?

[1 mark]

A Bromine has a higher electronegativity than iodine.

B Bromine has a lower boiling point than iodine.

C Bromine has a lower relative molecular mass than iodine.

D Bromine is a better oxidising agent than iodine.

1 3 Which equation represents a redox reaction?

[1 mark]

A $[\text{Co}(\text{H}_2\text{O})_6]^{2+} + 4\text{Cl}^- \rightarrow [\text{CoCl}_4]^{2-} + 6\text{H}_2\text{O}$

B $[\text{Fe}(\text{H}_2\text{O})_6]^{2+} + 2\text{OH}^- \rightarrow [\text{Fe}(\text{H}_2\text{O})_4(\text{OH})_2] + 2\text{H}_2\text{O}$

C $\text{MnO}_2 + 4\text{HCl} \rightarrow \text{MnCl}_2 + 2\text{H}_2\text{O} + \text{Cl}_2$

D $\text{CO}_2 + 2\text{NaOH} \rightarrow \text{Na}_2\text{CO}_3 + \text{H}_2\text{O}$

1 4 What is the oxidation state of N in N_2O_4 ?

[1 mark]

A -4

B -2

C +2

D +4

Turn over ►




1 5

Which compound does **not** form a racemic mixture when reacted with KCN, followed by dilute acid?

[1 mark]**A** Ethanal**B** Propanone**C** Butanone**D** Pentan-2-one**1 6**

The structure shows a section of a polymer chain formed from the random polymerisation of two different monomers.



Which pair of monomers could produce this polymer?


[1 mark]**A** $\text{CH}_2=\text{C}(\text{CH}_3)\text{Cl}$ and $\text{CHCl}=\text{CHCl}$ **B** $\text{CH}_2=\text{C}(\text{CH}_3)\text{Cl}$ and $(\text{CH}_3)_2\text{C}=\text{C}(\text{CH}_3)_2$ **C** $(\text{CH}_3)_2\text{C}=\text{CH}_2$ and $\text{CH}_2=\text{C}(\text{CH}_3)\text{Cl}$ **D** $(\text{CH}_3)_2\text{C}=\text{CH}_2$ and $\text{CHCl}=\text{C}(\text{CH}_3)\text{Cl}$ 

1 7


Which of the following curly arrows does **not** show the formation of a new covalent bond?

[1 mark]


A




B



C



D




Turn over for the next question

Turn over ►



1 8 The equation for the complete hydrogenation of buta-1,3-diene is



The table shows some mean bond enthalpies.

Bond	Mean bond enthalpy / kJ mol^{-1}
C–C	348
C=C	612
C–H	412
H–H	436

What is the enthalpy change, in kJ, for this reaction?

[1 mark]

A –684

B –248

C +140

D +576

1 9 Which Group 7 anion would give a white precipitate when dilute nitric acid is added, followed by a few drops of silver nitrate solution?

[1 mark]

A F^-

B Cl^-

C Br^-

D I^-




2 0 Which element could be sulfur?

[1 mark]

	Melting point of element	Solution of oxide	
A	high	acidic	<input type="checkbox"/>
B	high	alkaline	<input type="checkbox"/>
C	low	acidic	<input type="checkbox"/>
D	low	alkaline	<input type="checkbox"/>

2 1 The structure of aspartic acid is



Which structure is formed by aspartic acid in solution at pH 14?

[1 mark]

- A**
- $$\begin{array}{c} \text{H}_3\text{N}^+ - \text{CH} - \text{COO}^- \\ | \\ \text{H}_2\text{C} - \text{COO}^- \end{array}$$
-
- B**
- $$\begin{array}{c} \text{H}_2\text{N} - \text{CH} - \text{COO}^- \\ | \\ \text{H}_2\text{C} - \text{COO}^- \end{array}$$
-
- C**
- $$\begin{array}{c} \text{H}_2\text{N} - \text{CH} - \text{COO}^- \\ | \\ \text{H}_2\text{C} - \text{COOH} \end{array}$$
-
- D**
- $$\begin{array}{c} \text{H}_3\text{N}^+ - \text{CH} - \text{COOH} \\ | \\ \text{H}_2\text{C} - \text{COOH} \end{array}$$
-





Turn over ►



2	2
---	---

 Which structure shows an *E*- isomer?



[1 mark]



2 3 Benzene reacts with propanoyl chloride in the presence of an aluminium chloride catalyst.





Which compound is formed in this reaction?

[1 mark]



2 4 Which is a correct equation for a propagation step in the reaction of butane with chlorine to form dichlorobutane?

[1 mark]




Turn over for the next question

Turn over ►



2 5 The overall equation for the manufacture of sodium carbonate is



The relative formula masses of the reactants and products are

$$\text{CaCO}_3 = 100.1$$

$$\text{NaCl} = 58.5$$

$$\text{Na}_2\text{CO}_3 = 106.0$$

$$\text{CaCl}_2 = 111.1$$

What is the percentage atom economy of this process?

[1 mark]

A $\frac{106.0 \times 100}{100.1 + 58.5}$

B $\frac{106.0 \times 100}{106.0 + 111.1}$

C $\frac{111.1 \times 100}{100.1 + 58.5}$

D $\frac{111.1 \times 100}{100.1 + 106.0}$



2 6 Measurements made during an experiment in which compound **X** is dissolved in water are

mass of X	= 3.85 g
volume of solution formed	= 50.0 cm ³
initial temperature of water	= 18.7 °C
final temperature of solution	= 22.4 °C


The specific heat capacity of the solution = 4.18 J K⁻¹ g⁻¹
You should assume that the density of the solution is 1.00 g cm⁻³

What is the heat change, in joules, for this process?

[1 mark]

- A** $3.85 \times 4.18 \times (22.4 - 18.7)$
- B** $\frac{3.85 \times 4.18 \times (22.4 - 18.7)}{1000}$
- C** $50.0 \times 4.18 \times (22.4 - 18.7)$
- D** $\frac{50.0 \times 4.18 \times (22.4 - 18.7)}{1000}$

2 7 A student investigated the enthalpy change of the neutralisation reaction



The student mixed 20.0 cm³ of 1.85 mol dm⁻³ KOH(aq) with 30.0 cm³ of 1.75 mol dm⁻³ HCl(aq).

The heat change calculated was 1656 J

What is the enthalpy change, in kJ mol⁻¹, for this reaction?

[1 mark]

- A** - 44.8
- B** - 31.5
- C** + 31.5
- D** + 44.8

Turn over ►



2 8 The equation for the reaction between gaseous hydrogen and iodine is



An equilibrium mixture of gaseous hydrogen, iodine and hydrogen iodide was prepared at 500 K


The pressure and temperature of the equilibrium mixture were both increased.

Which statements describe the effects of these changes on the value of K_p ?

[1 mark]

	Effect of increase in pressure	Effect of increase in temperature	
A	K_p increases	K_p increases	<input type="checkbox"/>
B	K_p increases	K_p decreases	<input type="checkbox"/>
C	K_p unchanged	K_p increases	<input type="checkbox"/>
D	K_p unchanged	K_p decreases	<input type="checkbox"/>

2 9 The strengths of the weak acids 2-chloropropanoic acid and hydroxyethanoic acid are compared.



Which statement is correct?

[1 mark]

- A** Hydroxyethanoic acid is a stronger acid than 2-chloropropanoic acid.
- B** The pH of a $0.100 \text{ mol dm}^{-3}$ solution of hydroxyethanoic acid is 1.00
- C** The $\text{p}K_a$ value of 2-chloropropanoic acid is -2.83
- D** The K_a value of hydroxyethanoic acid is $1.48 \times 10^{-4} \text{ mol dm}^{-3}$




3 0 Which compound is butanoic anhydride?

[1 mark]



3 1 The equation for a reaction is



The rate of this reaction could be studied by following the

[1 mark]

A decrease in partial pressure of A

B increase in concentration of B

C decrease in concentration of C

D increase in partial pressure of D





Turn over for the next question

Turn over ►



3 2 Which pair of reagents would **not** react to give ethane-1,2-diol as one of the products?

[1 mark]




3 3


Which diagram shows the correct apparatus for the conversion of hexan-1-ol to hexanal?

[1 mark]


A




B



C



D



A

B

C

D

END OF QUESTIONS

30



There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**

Copyright information

For confidentiality purposes, acknowledgements of third party copyright material will be published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from www.oxfordqaexams.org.uk after the live examination series.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and Oxford International AQA Examinations will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2018 Oxford International AQA Examinations and its licensors. All rights reserved.

