

**INTERNATIONAL A-LEVEL
MATHEMATICS**

MA03

(9660/MA03) Unit P2 Pure Mathematics

Mark scheme

June 2024

Version: 1.0 Final



Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this mark scheme are available from oxfordaqa.com

Copyright information

OxfordAQA retains the copyright on all its publications. However, registered schools/colleges for OxfordAQA are permitted to copy material from this booklet for their own internal use, with the following important exception: OxfordAQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Copyright © 2024 OxfordAQA International Examinations and its licensors. All rights reserved.

Key to mark scheme abbreviations

M	Mark is for method
m	Mark is dependent on one or more M marks and is for method
A	Mark is dependent on M or m marks and is for accuracy
B	Mark is independent of M or m marks and is for method and accuracy
E	Mark is for explanation
√ or ft	Follow through from previous incorrect result
CAO	Correct answer only
CSO	Correct solution only
AWFW	Anything which falls within
AWRT	Anything which rounds to
ACF	Any correct form
AG	Answer given
SC	Special case
oe	Or equivalent
A2, 1	2 or 1 (or 0) accuracy marks
-x EE	Deduct x marks for each error
NMS	No method shown
PI	Possibly implied
SCA	Substantially correct approach
sf	Significant figure(s)
dp	Decimal place(s)
ISW	Ignore subsequent working

Q	Answer	Marks	Comments
1(a)	$[2\cos\theta + \sqrt{5}\sin\theta =]$ $R\cos\theta\cos\alpha + R\sin\theta\sin\alpha$ $R = 3$ $\alpha = 48^\circ$ $3\cos(\theta - 48^\circ)$	M1 B1 A1	PI Condone AWRT 48
		3	

Q	Answer	Marks	Comments
1(b)(i)	[Min value =] -3	B1ft	
		1	

Q	Answer	Marks	Comments
1(b)(ii)	228°	B1	Condone AWRT 228
		1	

	Question 1 Total	5	
--	-------------------------	----------	--

Q	Answer	Marks	Comments
2(a)	$0 \leq f(x) \leq \sqrt{18}$ or $[0, \sqrt{18}]$	B1	oe Accept f and y for f(x)
		1	

Q	Answer	Marks	Comments
2(b)(i)	$[fg(x) =] \sqrt{18 - 2\left(\frac{3}{2x-1}\right)^2}$	B1	oe ISW
		1	

Q	Answer	Marks	Comments
2(b)(ii)	$[0 \leq] \left(\frac{3}{2x-1}\right) \leq 3$ $x \geq 1$ only	M1 A1	or better
		2	

Q	Answer	Marks	Comments
2(b)(iii)	$\sqrt{18 - 2\left(\frac{3}{2x-1}\right)^2} = 4$ $2\left(\frac{3}{2x-1}\right)^2 = 2$ $x = 2$ and $x = -1$ (rejected)	M1 A1 A1	<i>their</i> $\sqrt{18 - 2\left(\frac{3}{2x-1}\right)^2} = 4$ PI by seeing either of $x = 2$ or $x = -1$ Only this value of x included in final answer
		3	

Q	Answer	Marks	Comments
3	$4x^2 + 12x + 9 = (2x + 3)^2 \text{ or } (2(x + 1.5))^2$ <p>or</p> $4x^2 + 12x + 9 = 4\left(x + \frac{3}{2}\right)^2$ <p>Translation $\begin{bmatrix} k \\ 0 \end{bmatrix}$</p> $\begin{bmatrix} -1.5 \\ 0 \end{bmatrix}$ <p>Stretch</p> <p>SF 4, parallel to y-axis</p> <p>OR</p> <p>Stretch</p> <p>SF 0.5, parallel to x-axis</p> <p>Translation $\begin{bmatrix} k \\ 0 \end{bmatrix}$</p> $\begin{bmatrix} -1.5 \\ 0 \end{bmatrix}$	<p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>[M1]</p> <p>[A1]</p> <p>[M1]</p> <p>[A1]</p>	<p>For either factorisation</p> <p>Translation $\begin{bmatrix} k \\ 0 \end{bmatrix}$</p> $\begin{bmatrix} -3 \\ 0 \end{bmatrix}$ <p>Stretch</p> <p>SF 0.5, parallel to x-axis</p> <p>Stretch</p> <p>SF 4, parallel to y-axis</p> <p>Translation $\begin{bmatrix} k \\ 0 \end{bmatrix}$</p> $\begin{bmatrix} -1.5 \\ 0 \end{bmatrix}$

	Question 3 Total	5	
--	-------------------------	----------	--

Q	Answer	Marks	Comments
4(a)	$X = Ae^{-kt}$		
	$\frac{dX}{dt} = -kAe^{-kt} = -kX$	B1	AG Must be convincingly shown
		1	

Q	Answer	Marks	Comments
4(b)	$X = Ae^{-kt}$		
	$t = 0, X = 90, A = 90$	B1	
	$t = 5, X = 80, 80 = 90e^{-5k}$	M1	
	$e^{-5k} = \frac{8}{9} \left[k = \frac{1}{5} \ln\left(\frac{9}{8}\right) = 0.02355\dots \right]$	A1	Allow $k = 0.024$ AWRT
	$X = 22.5$		
	$22.5 = 90e^{-kt}$		oe
	$t = -\frac{1}{k} \ln\left(\frac{22.5}{90}\right) \left[= -\frac{5}{\ln\left(\frac{9}{8}\right)} \ln\left(\frac{1}{4}\right) \right]$	m1	PI
	$t = 58.8$ [mins]	A1	Condone 58.7 to 58.9
		5	


	Question 4 Total	6	
--	-------------------------	----------	--

Q	Answer	Marks	Comments												
5(a)	<table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th>x</th> <th>y</th> </tr> </thead> <tbody> <tr> <td>0</td> <td>$3 - \tan^{-1}(-0.5) = 3.4636476$</td> </tr> <tr> <td>0.5</td> <td>$3 - \tan^{-1}(0.5) = 2.5363524$</td> </tr> <tr> <td>1</td> <td>$3 - \tan^{-1}(1.5) = 2.0172063$</td> </tr> <tr> <td>1.5</td> <td>$3 - \tan^{-1}(2.5) = 1.8097101$</td> </tr> <tr> <td>2</td> <td>$3 - \tan^{-1}(3.5) = 1.7075033$</td> </tr> </tbody> </table> $\frac{1}{3} \times 0.5(3.46\dots + 1.707\dots + 4(2.536\dots + 1.809\dots) + 2 \times 2.017\dots)$ $\int_0^2 3 - \tan^{-1}(2x - 0.5) dx = 4.432$	x	y	0	$3 - \tan^{-1}(-0.5) = 3.4636476$	0.5	$3 - \tan^{-1}(0.5) = 2.5363524$	1	$3 - \tan^{-1}(1.5) = 2.0172063$	1.5	$3 - \tan^{-1}(2.5) = 1.8097101$	2	$3 - \tan^{-1}(3.5) = 1.7075033$	<p>B1</p> <p>All five correct x values (and no extra used) PI by five correct y values</p> <p>M1</p> <p>At least four correct y values in exact form or decimals, rounded or truncated to two dp or better (in table or formula)</p> <p>m1</p> <p>Correct sub into formula with $h = 0.5$ oe and at least four correct y values either listed, with + signs, or totalled (PI by AWRT correct answer)</p> <p>A1</p> <p>CAO, must see this value exactly</p>	4
x	y														
0	$3 - \tan^{-1}(-0.5) = 3.4636476$														
0.5	$3 - \tan^{-1}(0.5) = 2.5363524$														
1	$3 - \tan^{-1}(1.5) = 2.0172063$														
1.5	$3 - \tan^{-1}(2.5) = 1.8097101$														
2	$3 - \tan^{-1}(3.5) = 1.7075033$														

Q	Answer	Marks	Comments
5(b)(i)	$f(x) = 0.5 \tan(3 - x) + 0.25 - x$ $f(1.7) = 0.35\dots$ $f(1.8) = -0.26\dots$ Change of sign, $1.7 < \alpha < 1.8$	<p>M1</p> <p>Both values rounded or truncated to at least 1 sf</p> <p>A1</p> <p>Must have both statement and interval in words or symbols or comparing 2 sides: M1 $y(1.7) = 2.05 > 1.7$ and $y(1.8) = 1.54 < 1.8$ A1: Conclusion as before</p>	2

Q	Answer	Marks	Comments
5(b)(ii)	$0.5 \tan(3-x) = x - 0.25$ $\tan(3-x) = 2x - 0.5$ $3-x = \tan^{-1}(2x-0.5)$ $x = 3 - \tan^{-1}(2x-0.5)$	M1 A1	oe AG Must be convincingly shown
		2	

Q	Answer	Marks	Comments
5(b)(iii)	$x_2 = 1.761$ $x_3 = 1.749$	B1 B1	
		2	

Q	Answer	Marks	Comments
5(b)(iv)	 <p>$y = x$</p> <p>$y = 3 - \tan^{-1}(2x - 0.5)$</p> <p>1.8</p> <p>1.6</p> <p>1.6</p> <p>x_1 x_3 x_2 1.8</p> <p>x_2 and x_3 in (approx.) correct position on axis</p>	<p>M1</p> <p>A1</p>	<p>Cobweb diagram</p>
		2	
	Question 5 Total	12	

Q	Answer	Marks	Comments
6(a)	<p>OR</p> $(\sin\theta - \operatorname{cosec}\theta)(\cos\theta - \sec\theta)$ $= \sin\theta\cos\theta - \frac{\sin\theta}{\cos\theta} - \frac{\cos\theta}{\sin\theta} + \frac{1}{\sin\theta\cos\theta}$ $= 0.5\sin 2\theta + \frac{-\sin^2\theta - \cos^2\theta + 1}{\sin\theta\cos\theta}$ $= 0.5\sin 2\theta + \frac{-1+1}{\sin\theta\cos\theta}$ $= 0.5\sin 2\theta$	<p>[B1]</p> <p>[M1]</p> <p>[A1]</p>	<p>All in terms of sin/cos</p> <p>Correct use of trig identity</p> <p>AG Must be convincingly shown</p>
		3	

Q	Answer	Marks	Comments
7(a)	$\left[\int \frac{x-2}{2x^2-8x+3} dx = \right]$ $a \ln(2x^2-8x+3) [+c]$ $= 0.25 \ln(2x^2-8x+3) [+c]$	<p>M1</p> <p>A1</p>	<p>ISW</p> <p>Accept $0.25 \ln(x^2-4x+1.5) [+c]$</p>
		2	

Q	Answer	Marks	Comments
7(b)	$\frac{5x+1}{5x-1} = \frac{5x-1+2}{5x-1} \left[= 1 + \frac{2}{5x-1} \right]$ $\left[\int \frac{5x+1}{5x-1} dx = \right] x + \frac{2}{5} \ln(5x-1) [+c]$	<p>B1</p> <p>M1 A1</p>	<p>PI</p> <p>M1: $x + k \ln(5x-1)$</p> <p>A1: ACF</p> <p>eg $x - \frac{1}{5} + \frac{2}{5} \ln(5x-1) [+c]$</p>
		3	

Q	Answer	Marks	Comments
7(c)	$\cos 6x = 1 - 2\sin^2 3x$ $\sin^2 3x = \frac{1}{2}(1 - \cos 6x)$ $\int \sin^2 3x dx = \frac{1}{2} \left(x - \frac{1}{6} \sin 6x \right) [+c]$	<p>B1</p> <p>M1 A1</p>	<p>M1: $ax + b \sin 6x$</p> <p>A1: ACF</p> <p>eg $\frac{x}{2} - \frac{\sin 3x \cos 3x}{6} [+c]$</p>
		3	

	Question 7 Total	8	
--	-------------------------	----------	--

Q	Answer	Marks	Comments
8(a)	$A(0, 5) \quad B(1.25, 0)$	B1	
		1	

Q	Answer	Marks	Comments
8(b)(i)	$y = (5 - 4x)e^{-0.5x}$ $\frac{dy}{dx} = (5 - 4x) \times -0.5e^{-0.5x} + (-4) \times e^{-0.5x}$ $\frac{dy}{dx} = (2x - 6.5)e^{-0.5x}$	M1 A1 A1	May use $y = 5e^{-0.5x} - 4xe^{-0.5x}$ M1: At least one exponential term differentiated correctly A1: All terms correct, unsimplified CAO (no ISW)
		3	

Q	Answer	Marks	Comments
8(b)(ii)	$(2x - 6.5)e^{-0.5x} = 0$ $2x - 6.5 = 0$ $x = 3.25$ $y = (5 - 4 \times 3.25)e^{-1.625}$ $y = -8e^{-1.625}$ $(3.25, -8e^{-1.625})$	M1 A1 A1	Sets <i>their</i> derivative equal to zero and attempts to solve ACF ISW
		3	

Q	Answer	Marks	Comments
8(c)	$\text{Area} = 0.5 \times 5 \times 1.25 - \int_0^{1.25} (5 - 4x)e^{-0.5x} dx$ $\int (5 - 4x)e^{-0.5x} dx$ $u = 5 - 4x \quad \frac{dv}{dx} = e^{-0.5x}$ $\frac{du}{dx} = -4 \quad v = -2e^{-0.5x}$ $\int (5 - 4x)e^{-0.5x} dx = (5 - 4x) \times (-2e^{-0.5x}) - \int -2e^{-0.5x} \times (-4) dx$ $= (5 - 4x)(-2e^{-0.5x}) + 16e^{-0.5x} \quad [+ c]$ $[\text{Area} = 0.5 \times 5 \times 1.25 -]$ $\left((0 + 16e^{-0.625}) - (-10 + 16) \right)$ $= 9.125 - 16e^{-0.625}$	<p>B1</p> <p>M1</p> <p>m1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>6</p>	<p>PI (by later work)</p> $\int 5e^{-0.5x} - 4xe^{-0.5x} dx$ $u = -4x \quad \frac{dv}{dx} = e^{-0.5x}$ $\frac{du}{dx} = -4 \quad v = -2e^{-0.5x}$ <p>Correct use of parts formula</p> $\int 5e^{-0.5x} - 4xe^{-0.5x} dx = -10e^{-0.5x} + (-4x) \times (-2e^{-0.5x}) - \int -2e^{-0.5x} \times (-4) dx$ $= -10e^{-0.5x} + 8xe^{-0.5x} + 16e^{-0.5x} \quad [+ c]$ <p>Correct substitution of limits into $(5 - 4x)(pe^{-0.5x}) + qe^{-0.5x}$ oe PI by</p> <p>ACF</p>

	Question 8 Total	13	
--	-------------------------	-----------	--

Q	Answer	Marks	Comments
9(c)	$\text{Area} = \frac{1}{2}(\pi a - 2a) \times a \left(\frac{\pi}{2} - 1 \right)$ $= \frac{1}{4} a^2 (\pi - 2)^2$	<p>M1</p> <p>A1</p>	<p>Uses <i>their</i> exact intercepts</p> <p>AG Must be convincingly shown</p>
		2	
	Question 9 Total	10	

Q	Answer	Marks	Comments
10(a)	$y = 4 \times 4 - (4 - y)^2$ $y = 16 - 16 + 8y - y^2$ $y^2 - 7y = 0 \Rightarrow y = 0, 7$ $AB = 7$	 M1 A1	 Attempt to solve for y PI
		2	

Q	Answer	Marks	Comments
10(b)	$\frac{dy}{dx} = 4 - 2(x - y) \left(1 - \frac{dy}{dx}\right)$ $\left[\frac{dy}{dx} = \frac{2x - 2y - 4}{2x - 2y - 1} \right]$ $\frac{dy}{dx} = 0, \quad x - y = 2$ $\left[\begin{aligned} (x - y)^2 &= 4x - y \\ &= (x - y) + 3x \end{aligned} \right]$ $2^2 = 2 + 3x$ $x = \frac{2}{3}$ $\frac{2}{3} - y = 2 \Rightarrow y = -\frac{4}{3}$ $\left(\frac{2}{3}, -\frac{4}{3} \right)$	 M1 A1 m1 A1	M1: Attempt at implicit differentiation $y \frac{dy}{dx}$ or $x \frac{dy}{dx}$ seen A1: All correct oe Substitutes <i>their</i> $ax + by = c$ in original equation oe $x = \frac{2}{3}, y = -\frac{4}{3}$
		4	

	Question 10 Total	6	
--	--------------------------	----------	--

Q	Answer	Marks	Comments
11(a)	$\frac{75}{(5-x)(5+2x)^2} = \frac{A}{5-x} + \frac{B}{5+2x} + \frac{C}{(5+2x)^2}$ $75 = A(5+2x)^2 + B(5-x)(5+2x) + C(5-x)$ $x=5: 75 = A(15)^2 \Rightarrow A = \frac{1}{3}$ $x=-2.5: 75 = C(7.5) \Rightarrow C = 10$ $x^2: 0 = 4A - 2B \Rightarrow B = \frac{2}{3}$ $\left[\frac{75}{(5-x)(5+2x)^2} = \right]$ $\frac{1/3}{5-x} + \frac{2/3}{5+2x} + \frac{10}{(5+2x)^2}$	<p>M1</p> <p>A1</p> <p>A1</p> <p>A1</p>	<p>Correctly eliminating fractions PI by at least one correct value for A, B or C</p> <p>At least one constant correct</p> <p>At least two constants correct</p> <p>All correct</p>
		4	

Q	Answer	Marks	Comments
11(b)	$(5+2x)^{-1} = 5^{-1} \left(1 + \frac{2x}{5} \right)^{-1}$ $= \frac{1}{5} \left(1 + (-1) \left(\frac{2x}{5} \right) + \frac{(-1) \times (-2) \times \left(\frac{2x}{5} \right)^2}{2} \right) [+...]$ $= \frac{1}{5} - \frac{2}{25}x + \frac{4}{125}x^2$	<p>M1</p> <p>A1</p>	<p>All correct</p> <p>Condone $\frac{1}{5} \left(1 - \frac{2}{5}x + \frac{4}{25}x^2 \right)$</p>
		2	

Q	Answer	Marks	Comments
11(c)	$(5-x)^{-1} = \frac{1}{5} \left(1 + \frac{1}{5}x + \frac{1}{25}x^2 \right)$	M1	Either expansion correct
	$(5+2x)^{-2} = \frac{1}{25} \left(1 - \frac{4}{5}x + \frac{12}{25}x^2 \right)$	A1	Both correct
	$f(x) =$		
	$\frac{1}{3} \left(\frac{1}{5} + \frac{1}{25}x + \frac{1}{125}x^2 \right) + \frac{2}{3} \left(\frac{1}{5} - \frac{2}{25}x + \frac{4}{125}x^2 \right)$	M1	Correct substitution of at least two of their expressions with their <i>A</i> , <i>B</i> , <i>C</i>
	$+ 10 \left(\frac{1}{25} - \frac{4}{125}x + \frac{12}{625}x^2 \right)$	A1ft	All correct
$= \frac{3}{5} - \frac{9}{25}x + \frac{27}{125}x^2$	A1		
		5	

	Question 11 Total	11	
--	--------------------------	-----------	--

Q	Answer	Marks	Comments
12	$x(16 - y^2) = 2(x^2 + 5) \frac{dy}{dx}$ $\int \frac{x}{x^2 + 5} dx = \int \frac{2}{16 - y^2} dy$ $\frac{2}{16 - y^2} = \frac{A}{4 + y} + \frac{B}{4 - y}$ $2 = A(4 - y) + B(4 + y)$ $A = \frac{1}{4} \quad B = \frac{1}{4}$ $\left[\int \frac{x}{x^2 + 5} dx = \int \frac{1/4}{4 + y} + \frac{1/4}{4 - y} dy \right]$ $\frac{1}{2} \ln(x^2 + 5) = \frac{1}{4} \ln(4 + y) - \frac{1}{4} \ln(4 - y) [+c]$ <p>At (0, 1) $2 \ln(5) = \ln\left(\frac{5}{3}\right) + c$</p> $c = \ln 15$ $2 \ln(x^2 + 5) = \ln(4 + y) - \ln(4 - y) + \ln 15$ $(x^2 + 5)^2 = \frac{15(4 + y)}{4 - y}$ $(x^2 + 5)^2 (4 - y) - 15y = 60$ $y = \frac{4(x^2 + 5)^2 - 60}{(x^2 + 5)^2 + 15}$	<p>M1</p> <p>m1</p> <p>A1</p> <p>m1 A1</p> <p>m1</p> <p>A1</p> <p>M1</p> <p>A1</p>	<p>Separates variables Condone omission of integral signs</p> <p>Attempt to use partial fractions PI by correct integration of $\int \frac{k}{16 - y^2} dy$</p> <p>Both correct PI by correct integration of $\int \frac{k}{16 - y^2} dy$</p> <p>m1: Attempt to integrate both sides A1: Both sides integrated correctly</p> <p>Attempt to find c, or to use limits for x and y</p> <p>or $c = \ln \frac{1}{15}$</p> <p>oe eg $\ln\left(\frac{(x^2 + 5)^2}{15}\right) = \ln\left(\frac{4 + y}{4 - y}\right)$</p> <p>Attempt to isolate y from $(x^2 + 5)^2 = \frac{k(4 \pm y)}{4 \pm y}$, $k \neq 0, 1$</p> <p>ACF eg $y = 4 - \frac{120}{x^4 + 10x^2 + 40}$</p>
	Question 12 Total	9	

Q	Answer	Marks	Comments
13(a)	$[AB: \mathbf{r}] = \begin{bmatrix} 2 \\ -2 \\ 6 \end{bmatrix} + \lambda \begin{bmatrix} 6 \\ 8 \\ -10 \end{bmatrix}$	B1	oe $\begin{bmatrix} 8 \\ 6 \\ -4 \end{bmatrix} + \lambda \begin{bmatrix} 6 \\ 8 \\ -10 \end{bmatrix}$
		1	

Q	Answer	Marks	Comments
13(b)(i)	$2 + 6\lambda = -2 + 4\mu$ or $8 + 6\lambda = -2 + 4\mu$ $-2 + 8\lambda = 3 - 5\mu$ or $6 + 8\lambda = 3 - 5\mu$ $\mu = 1, \lambda = 0$ or $\mu = 1, \lambda = -1$ $6 - 10\lambda = c + 3\mu$ or $-4 - 10\lambda = c + 3\mu$ $c = 3$	M1 A1 A1	Equating x and y coordinates AG Must be convincingly shown
		3	

Q	Answer	Marks	Comments
14(a)(i)	$\frac{dy}{dx} = \frac{\sin x(-\sin x) - \cos x(\cos x)}{\sin^2 x}$ $= \frac{-\sin^2 x - \cos^2 x}{\sin^2 x}$ $= -\operatorname{cosec}^2 x$	<p>M1</p> <p>A1</p>	<p>Attempt at quotient rule</p> <p>Must see a 'middle' line</p> <p>AG Must be convincingly shown</p>
		2	

Q	Answer	Marks	Comments
14(a)(ii)	$u = 1 + \cot x \Rightarrow \frac{du}{dx} = -\operatorname{cosec}^2 x$ $\int \frac{\operatorname{cosec}^2 x}{(1 + \cot x)^2} dx = -\int \frac{1}{u^2} du$ $= u^{-1} [+c]$ $\left[\frac{1}{u} \right]_{1+\sqrt{3}}^2 = \left[\frac{1}{1 + \cot x} \right]_{\frac{\pi}{6}}^{\frac{\pi}{4}}$ $\left[\int_{\frac{\pi}{6}}^{\frac{\pi}{4}} \frac{\operatorname{cosec}^2 x}{(1 + \cot x)^2} dx = \right] \frac{1}{2} - \frac{1}{1 + \sqrt{3}}$ $= 1 - \frac{\sqrt{3}}{2}$	<p>M1</p> <p>A1</p> <p>B1</p> <p>m1</p> <p>A1</p>	<p>Everything in terms of u</p> <p>Change limits or change back to x</p> <p>Substitute <i>their</i> limits into <i>their</i> expression of the form ku^{-1}</p>
		5	

