

**INTERNATIONAL AS
MATHEMATICS**

MA01

(9660/MA01) Unit P1 Pure Mathematics

Mark scheme

January 2025

Version: 1.0 Final



Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this mark scheme are available from www.oxfordaqa.com

Copyright information

OxfordAQA retains the copyright on all its publications. However, registered schools/colleges for OxfordAQA are permitted to copy material from this booklet for their own internal use, with the following important exception: OxfordAQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Copyright © 2025 OxfordAQA International Examinations and its licensors. All rights reserved.

Key to mark scheme abbreviations

M	Mark is for method
m	Mark is dependent on one or more M marks and is for method
A	Mark is dependent on M or m marks and is for accuracy
B	Mark is independent of M or m marks and is for method and accuracy
E	Mark is for explanation
√ or ft	Follow through from previous incorrect result
CAO	Correct answer only
CSO	Correct solution only
AWFW	Anything which falls within
AWRT	Anything which rounds to
ACF	Any correct form
AG	Answer given
SC	Special case
oe	Or equivalent
A2, 1	2 or 1 (or 0) accuracy marks
-x EE	Deduct x marks for each error
NMS	No method shown
PI	Possibly implied
SCA	Substantially correct approach
sf	Significant figure(s)
dp	Decimal place(s)
ISW	Ignore subsequent working

Q	Answer	Marks	Comments
1(a)(i)	$\frac{64}{125}$	B1	
		1	

Q	Answer	Marks	Comments
1(a)(ii)	-3	B1	
		1	

Q	Answer	Marks	Comments
2(a)	$[8 \times 8 - 5 \times 10 = 14 \Rightarrow] \quad 64 - 50 = 14$ or $\left[\frac{8}{5} \times 8 - \frac{14}{5} \Rightarrow \right] \quad \frac{64}{5} - \frac{14}{5} = 10$ or $[8 \times 8 - 5y = 14 \Rightarrow]$ $64 - 5y = 14$ $\Rightarrow y = 10$ and Hence l_1 passes through the point $A(8,10)$	B1	Either: Substitutes the coordinates of A into the LHS of the equation of l_1 and concludes it is equal to 14. Must see $64 - 50 = 14$ or Rearranges to $y = \frac{8}{5}x - \frac{14}{5}$ and substitutes into RHS concluding $y = 10$ Must see $\frac{64}{5} - \frac{14}{5} = 10$ or Substitutes $x = 8$ into LHS and solves for y . Must see $64 - 5y = 14$ before $y = 10$ Must have a concluding statement
		1	

Q	Answer	Marks	Comments
2(b)	$[8x - 5y = 14 \text{ and } 7x + 2y = -26 \Rightarrow]$ $(-2, -6)$ $[AB =] \sqrt{(8 - (-2))^2 + (10 - (-6))^2}$ $2\sqrt{89} \text{ or } \sqrt{356}$	B1 M1 A1	PI Solves equations simultaneously to find the coordinates of B Condone not given as coordinates but must be clearly identified. oe ft their coordinates of B CAO
		3	

Q	Answer	Marks	Comments
2(c)(i)	$[\text{Gradient of } l_1 =] \frac{8}{5}$	B1	oe
	$\frac{(k+7)-5}{3-(-2)} \left[= \frac{8}{5} \right]$	M1	oe Correct method for finding the gradient of l_3 in terms of k
	$\left[\frac{k+2}{5} = \frac{8}{5} \Rightarrow \right] k = 6$	A1	CAO
		3	

Q	Answer	Marks	Comments
2(c)(ii)	$y-5 = \frac{8}{5}(x-(-2))$	M1	oe Forms a correct equation for l_3 but not in the required form
	or $y-13 = \frac{8}{5}(x-3)$		May see $y = \frac{8}{5}x + c$ or $8x - 5y = c$ and substitution of coordinates of P or Q to find c but must be a complete method
	or $y = \frac{8}{5}x + \frac{41}{5}$		ft their gradient of l_1 from part (c)(i) ft their k from part (c)(i)
	$8x - 5y + 41 = 0$	A1	CAO Any integer multiple but must be in the correct form
		2	

	Question 2 Total	9	
--	-------------------------	----------	--

Q	Answer	Marks	Comments
3(a)	Arithmetic	E1	Any statement implying a common difference between consecutive terms
	The amount saved increases by the same amount each month	E1	
		2	


Q	Answer	Marks	Comments
3(b)	$[a = 240, d = 8 \text{ and } n = 22 \Rightarrow]$ $240 + (22 - 1) \times 8$	M1	PI oe Correct use of the formula for the n th term of an arithmetic series with values substituted
	[\$] 408	A1	CAO
		2	

Q	Answer	Marks	Comments
3(c)	$[a = 240, d = 8 \text{ and } n = 36 \Rightarrow]$ $\frac{1}{2} \times 36(2 \times 240 + (36 - 1) \times 8)$	M1	PI oe Correct use of the formula for the sum of the first n terms of an arithmetic series with values substituted
	[\$] 13 680	A1	CAO
		2	

Q	Answer	Marks	Comments
4(a)	$(-3)^3 - 4(-3) + 15 [= 0]$ $-27 + 12 + 15 = 0$	M1	Factor Theorem used with $x = -3$ substituted
		A1	oe Powers and products evaluated before being set equal to zero Factor Theorem not used scores M0 A0
		2	

Q	Answer	Marks	Comments
4(b)	$(x+3)(x^2 - 3x + 5)$	M1	b or c correct
		A1	CAO
		2	

Q	Answer	Marks	Comments
4(c)	$[b^2 - 4ac =] (-3)^2 - 4 \times 1 \times 5 [= -11]$ $-11 < 0$ therefore the equation $f(x) = 0$ has exactly one real root	M1	oe Correct attempt to evaluate the discriminant of $x^2 - 3x + 5$ ft Their answer to part (b) PI by -11 or a correct completed-square form for the quadratic factor
		A1ft	Their discriminant evaluated correctly with an indication that it is negative and a final conclusion or an indication that the minimum value of their correct completed-square form is positive and a final conclusion
		2	

Q	Answer	Marks	Comments
4(d)		<p>B1</p> <p>B1</p> <p>B1</p>	<p>Cubic curve of the correct form with minimum in the first quadrant and maximum in the second quadrant</p> <p>Correct value for y-intercept provided a graph is drawn Allow given as coordinates</p> <p>Correct value for x-intercept and no others provided a graph is drawn Allow given as coordinates</p>
		3	
	Question 4 Total	9	

Q	Answer	Marks	Comments
5(a)	$\frac{7x\sqrt{x} + 14x - 3\sqrt{x} - 6}{\sqrt{x} + 2} \times \frac{\sqrt{x} - 2}{\sqrt{x} - 2}$ $x - 4$ $7x^2 + 14x\sqrt{x} - 3x - 6\sqrt{x} - 14x\sqrt{x} - 28x + 6\sqrt{x} + 12$ $7x^2 - 31x + 12$ $\left[\frac{7x^2 - 31x + 12}{x - 4} = \right]$ $\frac{(7x - 3)(x - 4)}{x - 4}$ and $7x - 3$	<p>M1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>5</p>	<p>PI Intention to multiply numerator and denominator by $\sqrt{x} - 2$</p> <p>Correct denominator Must be seen as the denominator</p> <p>Unsimplified expression for the numerator Allow one error</p> <p>Correct simplified numerator</p> <p>Correct algebraic fraction with numerator factorised before correct answer in given form. If numerator not factorised allow evidence of correct algebraic division.</p> <p>SC2 for final answer of $7x - 3$ for methods other than rationalising the denominator</p>

Q	Answer	Marks	Comments
6(a)	$[(1+3x)^9 =]$ $[(1)^9 + 9(1)^8(3x) +]$ $36(1)^7(3x)^2 + 84(1)^6(3x)^3]$ $[a =] 324$ $[b =] 2268$	<p>M1</p> <p>For either $[1, 9], 36, 84$ oe unsimplified</p> <p>or $\binom{9}{2}(1)^7(3x)^2$ or $\binom{9}{3}(1)^6(3x)^3$</p> <p>oe, x not needed PI</p> <p>A1 Condone $324x^2$</p> <p>A1 Condone $2268x^3$</p>	<p>3</p>

Q	Answer	Marks	Comments
6(b)	$1+3x = \frac{17}{20}$ $[x =] -\frac{1}{20}$ $\left[\left(\frac{17}{20}\right)^9 \approx\right]$ $1+27\left(-\frac{1}{20}\right) + 324\left(-\frac{1}{20}\right)^2 + 2268\left(-\frac{1}{20}\right)^3$ $1 - \frac{27}{20} + \frac{81}{100} - \frac{567}{2000}$ and $\frac{353}{2000}$	<p>M1 PI by $x = -\frac{1}{20}$ seen substituted</p> <p>Method for finding correct value of x</p> <p>A1 Possibly seen embedded in later working.</p> <p>m1 oe Substitutes their $x = -\frac{1}{20}$ into their expansion in part (a)</p> <p>A1 oe Extra line of working simplifying powers and products before AG</p>	<p>4</p>

Q	Answer	Marks	Comments
7(a)(i)	$\left[y = ax^3 + bx^2 + cx^{-\frac{3}{2}} \Rightarrow \right]$ $\left[\frac{dy}{dx} = \right] 3ax^2 + 2bx - \frac{3}{2}cx^{-\frac{5}{2}}$	M1 A1	M1: oe At least two correct terms A1: oe Correct first derivative
		2	

Q	Answer	Marks	Comments
7(a)(ii)	$\left[\frac{d^2y}{dx^2} = \right] 6ax + 2b + \frac{15}{4}cx^{-\frac{7}{2}}$	M1 A1ft	oe At least two correct terms ft Through their first derivative oe Correct second derivative ft Their answer to part (a)(i) provided it contains a fractional power of x
		2	

Q	Answer	Marks	Comments
7(b)	Since a, b and c [and x] are positive then $\frac{d^2y}{dx^2} > 0$ Hence P is a minimum point	E1ft E1	States a, b and c [and x] are positive and therefore the second derivative is positive ft Their answer to part (a)(ii) provided all terms are positive Correct conclusion E0 E1 not possible
		2	

Q	Answer	Marks	Comments
8(a)	$\left[y = \frac{3}{4}x^2 - 12x + 21 \Rightarrow \right]$ $\left[\frac{dy}{dx} = \right] \frac{3}{2}x - 12$ $\left[x = 6 \Rightarrow \frac{dy}{dx} = \frac{3}{2} \times 6 - 12 = \right] -3$ $\left[m' \times (-3) = -1 \Rightarrow \right]$ $\left[m' = \right] \frac{1}{3}$ or $(-3) \times \frac{3}{2}$ or $-\frac{9}{2}$ $\frac{1}{3} \neq \frac{3}{2}$ or $(-3) \times \frac{3}{2} \neq -1$ or $-\frac{9}{2} \neq -1$ or $-3 \neq -\frac{2}{3}$ therefore l is not the normal to C at P	<p>M1</p> <p>A1</p> <p>B1ft</p> <p>E1ft</p>	<p>oe Correct derivative simplified or unsimplified.</p> <p>PI by correct gradient of tangent to C at P</p> <p>Correct gradient of tangent to C at P</p> <p>Correct gradient of normal to C at P or Correct expression for, or value of, the product of the gradient of tangent to C at P and the gradient of l</p> <p>ft Their gradient of tangent to C at P</p> <p>Compares gradient of normal to C at P to gradient of l and makes correct conclusion.</p> <p>or</p> <p>Compares $-\frac{9}{2}$ or $(-3) \times \frac{3}{2}$ to -1 and makes correct conclusion.</p> <p>ft Their gradient of normal to C at P provided it is not $\frac{3}{2}$</p>
		4	

Q	Answer	Marks	Comments
8(b)(i)	$\frac{3}{4}x^2 - 12x + 21 = \frac{3}{2}x - 33$	M1	oe Equates the equation of C with the equation of l
	$\frac{3}{4}x^2 - \frac{27}{2}x + 54 = 0$ or $3x^2 - 54x + 216 = 0$ and $x^2 - 18x + 72 = 0$	A1	oe Extra line of working showing a simplified quadratic equation set equal to zero before AG Must be convincingly shown
		2	

Q	Answer	Marks	Comments
8(b)(ii)	$[x^2 - 18x + 72 = 0 \Rightarrow]$	M1	Correct x-coordinate of Q Ignore $x = 6$ if seen as well
	$[x =] 12$ $(12, -15)$	A1	Correct coordinates of Q Condone not given as coordinates but must be clearly identified
		2	

Q	Answer	Marks	Comments
8(c)	$\left[\int \left(\frac{3}{4}x^2 - 12x + 21 \right) dx = \right]$		
	$\frac{1}{4}x^3 - 6x^2 + 21x [+c]$	M1 A1	M1: At least two correct terms simplified or unsimplified A1: Correct integration simplified or unsimplified
		2	

Q	Answer	Marks	Comments
8(d)	$\left[\int_{12}^{14} \left(\frac{3}{4}x^2 - 12x + 21 \right) dx = \right]$ $\left(\frac{1}{4}(14)^3 - 6(14)^2 + 21(14) \right)$ $- \left(\frac{1}{4}(12)^3 - 6(12)^2 + 21(12) \right)$ $\left[\int_{12}^{14} \left(\frac{3}{4}x^2 - 12x + 21 \right) dx = \right] -16$ $\frac{1}{2} \times (22 - 12) \times 15 [= 75]$ $[75 - 16 =] 59$	<p>M1</p> <p>A1</p> <p>M1</p> <p>A1ft</p>	<p>oe Correct attempt to evaluate the definite integral ft Their x-coordinate of Q PI by the correct value of the integral.</p> <p>CAO</p> <p>Correct method for finding the area of a relevant triangle ft Their coordinates of Q ft Their value for the definite integral provided it was negative and their value for the area of a relevant triangle provided both method marks awarded</p>
		4	
	Question 8 Total	14	

Q	Answer	Marks	Comments
9(a)	$ar = -48 \text{ or } \frac{a}{1-r} = 200$ $\left[a = -\frac{48}{r} \text{ and } a = 200(1-r) \Rightarrow \right]$ $-\frac{48}{r} = 200(1-r)$ $200r^2 - 200r - 48 = 0$ or $25r^2 - 25r - 6 = 0$ $r = -\frac{1}{5} \text{ and } r = \frac{6}{5}$ and <p>[since the series has a sum to infinity then]</p> $r = \frac{6}{5} \text{ is rejected}$ $[a =] 240$	<p>B1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>B1</p>	<p>oe Correct equation in terms of a and r using the second term or the sum to infinity</p> <p>oe Starts to solve the equations simultaneously and forms a single equation in r</p> <p>oe Forms a correct quadratic equation in r set equal to zero</p> <p>oe Finds both possible values of r and rejects $r = \frac{6}{5}$</p> <p>CAO</p>
		5	

Q	Answer	Marks	Comments
9(a) ALT	$ar = -48 \text{ or } \frac{a}{1-r} = 200$ $\left[r = -\frac{48}{a} \Rightarrow \right]$ $\frac{a}{1 - \left(-\frac{48}{a}\right)} = 200 \text{ or } \frac{a^2}{a+48} = 200$ $a^2 - 200a - 9600 = 0$ $[a = -40 \text{ or } 240]$ <p>If $a = -40$ then $r = \frac{6}{5}$</p> <p>If $a = 240$ then $r = -\frac{1}{5}$</p> <p>and</p> <p>[since the series has a sum to infinity then]</p> $r = \frac{6}{5} \text{ is rejected}$ $[a =] 240$	<p>B1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>B1</p>	<p>oe Correct equation in terms of a and r using the second term or the sum to infinity</p> <p>oe Starts to solve the equations simultaneously and forms a single equation in a</p> <p>oe Forms a correct quadratic equation in a set equal to zero</p> <p>oe Finds the correct corresponding value of r for each value of a and rejects $r = \frac{6}{5}$</p> <p>CAO</p>
		<p>5</p>	

Q	Answer	Marks	Comments
9(b)	$\left[\sum_{n=1}^{2k} \frac{625}{8} u_n = \right] \left[\frac{625}{8} \times \frac{240 \left(1 - \left(-\frac{1}{5} \right)^{2k} \right)}{1 - \left(-\frac{1}{5} \right)} \right]$ $\left[\sum_{n=1}^{2k} \frac{625}{8} u_n = \right] 5^6 (1 - 5^{-2k})$	<p>M1</p> <p>m1</p> <p>A1</p>	<p>oe Substitutes $r = -\frac{1}{5}$ and their $a = 240$ into the formula for the sum of the first $2k$ terms of the series</p> <p>In the correct form with the correct values of b and c or b and d or $b = \frac{1}{5}$, $c = -6$ and $d = 2$</p> <p>Condone $(-5)^{-2k}$ in place of 5^{-2k}</p> <p>CAO Correct final answer in the correct form</p>
		3	
	Question 9 Total	8	