

Please write clearly in block capitals.

Centre number

Candidate number

Surname

Forename(s)

Candidate signature

I declare this is my own work.

INTERNATIONAL AS

MATHEMATICS

(9660/MA01) Unit P1 Pure Mathematics

Wednesday 15 May 2024

07:00 GMT

Time allowed: 1 hour 30 minutes

Materials

- For this paper you must have the OxfordAQA Booklet of Formulae and Statistical Tables (enclosed).
- You may use a graphical calculator.

Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer the questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do all rough work in this book. Cross through any work you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 80.

Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- Show all necessary working; otherwise marks may be lost.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
TOTAL	



Answer **all** questions in the spaces provided.

- 1 (a)** The n th term of a sequence is x_n where

$$x_{n+1} = x_n + 5$$

The fourth term $x_4 = 30$

- 1 (a) (i)** Find the value of x_1

Circle your answer.

[1 mark]

5

10

15

20

- 1 (a) (ii)** Find the value of n for which $x_n = 60$

Circle your answer.

[1 mark]

8

9

10

11



1 (b) Two sequences are generated by the formulae

$$u_{n+1} = pu_n + 12 \quad \text{and} \quad t_{n+1} = 2pt_n + 3$$

where p is a constant.

The sequences have the same non-zero limit L as $n \rightarrow \infty$

Find the value of L and the value of p

[4 marks]

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.
$$L = \quad \quad \quad p =$$

6

Turn over ►



2 (a) Given that $x \neq 0$ and $y \neq 0$ simplify

$$\frac{\left(3x^7y^{\frac{2}{3}}\right)^6}{\sqrt{9x^4y^{\frac{3}{2}}}}$$

Give your answer in the form

$$ax^by^c$$

where a , b and c are constants.

[3 marks]

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Answer



2 (b)

can be written in the form $d + e\sqrt{w}$, where d and e are integers.

[3 marks]


[illegible]

Turn over for the next question

Turn over ►



- 3** The diagram shows the lines l_1 and l_2 and the points $P(6, -4)$ and $Q(-1, -3)$



The line l_1 passes through the point P and intersects the line l_2 at the point Q

The lines l_1 and l_2 are perpendicular.

- 3 (a)** Show that the equation of l_2 is

$$y = 7x + 4$$

[3 marks]



3 (b) The point R lies on l_2

The x -coordinate of R is a where $a > 0$

The distance QR is $6\sqrt{10a}$

3 (b) (i) Show that

$$5a^2 - 26a + 5 = 0$$

[3 marks]

3 (b) (ii) Hence find the coordinates of the two possible positions of R

[3 marks]

Answer _____ and _____



- 4** The third, fourth and sixth terms of a finite arithmetic series are

$$u_3 = p - 4, \quad u_4 = 2p - 18 \quad \text{and} \quad u_6 = 5p - 67$$

where p is a constant.

- 4 (a)** Show that $p = 21$

[2 marks]

- 4 (b)** The last term of the series is $u_m = 990$

Find the value of m

[4 marks]

$m =$ _____



4 (c) The sum of all the **even** terms of the series is S

Find the value of S

[3 marks]

$S =$ _____

9

Turn over for the next question

Turn over ►



- 5** The function f is given by

$$f(x) = 3x^2 + (4k + 5)x + 10k + a$$

where k and a are constants.

When $f(x)$ is divided by $(x + 5)$ the remainder is $59 - 17k$

- 5 (a)** Use the Remainder Theorem to show that

$$a = 9 - 7k$$

[2 marks]

- 5 (b)** The curve C has equation $y = f(x)$

The line L has gradient 7 and intersects the y -axis at the point $(0, -3k)$

The curve C and the line L intersect at two distinct points.

Find the possible values of k

[8 marks]



[illegible]

Answer _____

Turn over ►



6 The curve C has equation

$$y = 6x^2 + p$$

where p is a constant.

6 (a) Write down the coordinates of the vertex of C

[1 mark]

Answer

6 (b) The curve D has equation

$$y = 6x^2 - 48x + 103$$

By completing the square describe the single transformation which maps C onto D

[5 marks]



7 (a) Show that

$$\left(12x^{\frac{4}{5}} + 1\right)\left(x^{-\frac{4}{5}} - 12\right) = x^{-\frac{4}{5}} - 144x^{\frac{4}{5}}$$

[1 mark]

7 (b) Hence find $\int \left(12x^{\frac{4}{5}} + 1\right)\left(x^{-\frac{4}{5}} - 12\right) dx$

Give your answer in the form

$$qx^r(1 - sx^t) + c$$

where q , r , s and t are constants to be found, and c is the constant of integration.

[3 marks]

Answer _____

Turn over ►



8

$$f(x) = \frac{k - 6\sqrt{x} + 5x}{\sqrt{x}} + \frac{9}{2}$$

where $x > 0$ and k is a constant.

The point P has x -coordinate $\frac{1}{4}$ and lies on C

8 (a)

$$5 - 4k$$

[4 marks]

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins or other markings on the paper.

8 (b) The tangent to C at P is parallel to the line with equation $y = -43x + 9$

8 (b) (i) Show that $k = 12$

[1 mark]

8 (b) (ii) Find the equation of the tangent to C at P

Give your answer in the form $ax + by = c$ where a , b and c are integers.

[3 marks]

Answer _____

Question 8 continues on the next page

Turn over ►



8 (c) The curve C has one stationary point at Q

The x -coordinate of Q is $\frac{12}{5}$

Use the second derivative to determine whether Q is a maximum point or a minimum point.

[3 marks]

8 (d) State the values of x for which f is an increasing function.

[1 mark]

Answer _____



- 9** A student is testing the accuracy of the trapezium rule by using it to estimate the value of the integral I where

$$I = \int_{0.5}^{1.5} \left(\frac{1}{4x^2} + \frac{x^3}{6} \right) dx$$

- 9 (a) (i)** Complete the table giving your values to four decimal places.

[2 marks]

x	0.5	0.75	1	1.25	1.5
$\frac{1}{4x^2} + \frac{x^3}{6}$		0.5148	0.4167		

- 9 (a) (ii)** Hence use the trapezium rule with five ordinates (four strips) to find an estimate for I

Give your answer to three decimal places.

[3 marks]

Answer _____

Question 9 continues on the next page

Turn over ►



9 (b) (i) Find $\int \left(\frac{1}{4x^2} + \frac{x^3}{6} \right) dx$

[2 marks]

Answer _____

9 (b) (ii) Hence find the exact value of $\int_{0.5}^{1.5} \left(\frac{1}{4x^2} + \frac{x^3}{6} \right) dx$

[2 marks]

Answer _____



- Use your answers to **parts (a)(ii)** and **(b)(ii)** to determine whether the student is correct.

[illegible]

- [1 mark]**

Turn over for the next question

The diagram shows the first four squares of a sequence consisting of an infinite number of squares.

Diagram illustrating the sequence of squares with side lengths x cm, x^2 cm, x^3 cm, and x^4 cm, followed by an ellipsis indicating the sequence continues.

The area, x^2 , of the largest square is the first term.

The sum to infinity of the **perimeters** of the squares is P cm

[6 marks]

[illegible]

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

$P =$ _____

6

END OF QUESTIONS



There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**



*Do not write
outside the
box*

[illegible]

[illegible]