

Please write clearly in block capitals.

Centre number

Candidate number

Surname \_\_\_\_\_

Forename(s) \_\_\_\_\_

Candidate signature \_\_\_\_\_

I declare this is my own work.

# INTERNATIONAL A-LEVEL MATHEMATICS

(9660/MA04) Unit S2 Statistics

Tuesday 16 January 2024 07:00 GMT Time allowed: 1 hour 30 minutes

## Materials

- For this paper you must have the OxfordAQA Booklet of Formulae and Statistical Tables (enclosed).
- You may use a graphical calculator.

## Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer the questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do all rough work in this book. Cross through any work you do not want to be marked.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
<b>TOTAL</b>	

## Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 80.

## Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- Show all necessary working; otherwise marks may be lost.



Answer **all** questions in the spaces provided.

- 1** The probability density function  $f$  for a continuous random variable  $X$  is given by

$$f(x) = \begin{cases} \frac{1}{7}e^{-\frac{1}{7}x} & x \geq 0 \\ 0 & \text{otherwise} \end{cases}$$

- 1 (a)** State the name of the distribution of  $X$

[1 mark]

\_\_\_\_\_

Answer \_\_\_\_\_

- 1 (b)** Find the mean of  $X$

[1 mark]

\_\_\_\_\_

Answer \_\_\_\_\_

- 1 (c)** Find  $P(2 < X < 7)$

Give your answer to four significant figures.

[2 marks]

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

Answer \_\_\_\_\_



1 (d) It is given that  $P(X < a) = 0.8$

Find the value of  $a$

Give your answer to four significant figures.

[2 marks]

---

---

---

---

Answer \_\_\_\_\_

1 (e) Find  $P(X > 8 \mid X > 5)$

Give your answer to four significant figures.

[2 marks]

---

---

---

---

Answer \_\_\_\_\_

8
---

Turn over for the next question

Turn over ►







- 3 The continuous random variable  $X$  has probability density function  $f(x)$  defined by

$$f(x) = \begin{cases} \frac{1}{8}x^2 & 0 \leq x < 1 \\ k(x-1) + \frac{1}{8} & 1 \leq x \leq 6 \\ 0 & \text{otherwise} \end{cases}$$

where  $k$  is a constant.

- 3 (a) Find  $P(X < 1)$

Give your answer in exact form.

[2 marks]

---



---



---



---

Answer \_\_\_\_\_

- 3 (b) Show that  $k = \frac{2}{75}$

[3 marks]

---



---



---



---



---



---



---




---



3 (c) Draw the graph of  $y = f(x)$  for  $0 \leq x \leq 8$

[3 marks]



Question 3 continues on the next page

Turn over ►





- 4 (a)** The random variables  $C$ ,  $D$ ,  $E$  and  $F$  are all binomially distributed, as shown in the table.

Distribution
$C \sim B(10, 0.04)$
$D \sim B(200, 0.51)$
$E \sim B(400, 0.91)$
$F \sim B(100, 0.05)$

State, with a reason, which one of these distributions would be most suitable to approximate by a Poisson distribution.

**[2 marks]**

Distribution \_\_\_\_\_

Reason \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

**Question 4 continues on the next page**

**Turn over ►**





- 4 (c)** A different Poisson distribution is used to model the number of helicopters flying over the island in any given hour. The mean of this distribution is 2.7

Assume that the number of helicopters and the number of aeroplanes flying over the island in any given hour are independent random variables.

The random variable  $G$  represents the total number of helicopters and aeroplanes flying over the island in any given hour.

- 4 (c) (i)** Write down the parameter of  $G$

**[1 mark]**

---

Answer \_\_\_\_\_

- 4 (c) (ii)** It is given that  $P(G < a) > 0.95$  where  $a$  is an integer.

By considering probabilities, find the smallest possible value of  $a$

**[3 marks]**

---



---



---



---



---



---



---



---



---



---

Answer \_\_\_\_\_







- 6** The mass  $V$  grams of flour in a bag can be modelled by a normal distribution with mean 502 grams and standard deviation 2.7 grams.

A bakery purchases 30 bags of flour.

The masses of the bags are independent and each bag is selected at random.

- 6 (a)** Find the probability that the mean mass of the 30 bags is less than 501 grams.

Give your answer to four decimal places.

**[4 marks]**

---

---

---

---

---

---

---

---

---

---

Answer \_\_\_\_\_

- 6 (b)** Find the probability that the mass of flour in each of the 30 bags is more than 496 grams.

Give your answer to three decimal places.

**[4 marks]**

---

---

---

---

---

---

---

---

---

---

Answer \_\_\_\_\_











*Do not write  
outside the  
box*

---

---

---

---

---

---

---

      
**6**

**END OF QUESTIONS**



**There are no questions printed on this page**

*Do not write  
outside the  
box*

**DO NOT WRITE ON THIS PAGE  
ANSWER IN THE SPACES PROVIDED**









**There are no questions printed on this page**

*Do not write  
outside the  
box*

**DO NOT WRITE ON THIS PAGE  
ANSWER IN THE SPACES PROVIDED**

**Copyright information**

For confidentiality purposes, all acknowledgements of third-party copyright material are published in a separate booklet. This booklet is published after each live examination series and is available for free download from [www.oxfordaqa.com](http://www.oxfordaqa.com)

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and OxfordAQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team.

Copyright © 2024 OxfordAQA International Examinations and its licensors. All rights reserved.

