Cambridge International A Level

Published

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptions for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1: Marks must be awarded in line with: the specific content of the mark scheme or the generic level descriptors for the question the specific skills defined in the mark scheme or in the generic level descriptors for the question the standard of response required by a candidate as exemplified by the standardisation scripts.
GENERIC MARKING PRINCIPLE 2: Marks awarded are always **whole marks** (not half marks, or other fractions).
GENERIC MARKING PRINCIPLE 3: Marks must be awarded **positively**: marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate marks are awarded when candidates clearly demonstrate what they know and can do marks are not deducted for errors marks are not deducted for omissions answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.
GENERIC MARKING PRINCIPLE 4: Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.
GENERIC MARKING PRINCIPLE 5: Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).
GENERIC MARKING PRINCIPLE 6: Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Mathematics-Specific Marking Principles

1   Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing.

2   Unless specified in the question, non-integer answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected.

3   Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.

4   Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw).

5   Where a candidate has misread a number or sign in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 A or B mark for the misread.

6   Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.

Annotations guidance for centres

Examiners use a system of annotations as a shorthand for communicating their marking decisions to one another. Examiners are trained during the standardisation process on how and when to use annotations. The purpose of annotations is to inform the standardisation and monitoring processes and guide the supervising examiners when they are checking the work of examiners within their team. The meaning of annotations and how they are used is specific to each component and is understood by all examiners who mark the component.

We publish annotations in our mark schemes to help centres understand the annotations they may see on copies of scripts. Note that there may not be a direct correlation between the number of annotations on a script and the mark awarded. Similarly, the use of an annotation may not be an indication of the quality of the response.

The annotations listed below were available to examiners marking this component in this series.

Annotations

Annotation	Meaning
⌃	More information required
A0	Accuracy mark awarded zero
A1	Accuracy mark awarded one
B0	Independent accuracy mark awarded zero
B1	Independent accuracy mark awarded one
B2	Independent accuracy mark awarded two
BOD	Benefit of the doubt
BP	Blank Page
✖	Incorrect
Dep	Used to indicate DM0 or DM1

Annotation	Meaning
DM1	Dependent on the previous M1 mark(s)
FT	Follow through
~~	Indicate working that is right or wrong
Highlighter	Highlight a key point in the working
ISW	Ignore subsequent work
J	Judgement
JU	Judgement
M0	Method mark awarded zero
M1	Method mark awarded one
M2	Method mark awarded two
MR	Misread
O	Omission or Other solution
Off-page comment	Allows comments to be entered at the bottom of the RM marking window and then displayed when the associated question item is navigated to.
On-page comment	Allows comments to be entered in speech bubbles on the candidate response.
PE	Judgment made by the PE
Pre	Premature approximation
SC	Special case
SEEN	Indicates that work/page has been seen

Annotation	Meaning
SF	Error in number of significant figures
✔	Correct
TE	Transcription error
XP	Correct answer from incorrect working

PUBLISHED

Mark Scheme Notes

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

Types of mark

M   Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

A   Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

B   Mark for a correct result or statement independent of method marks.

DM or **DB**   When a part of a question has two or more 'method' steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly, when there are several B marks allocated. The notation DM or DB is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.

FT   Implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only.

A or B marks are given for correct work only (not for results obtained from incorrect working) unless follow through is allowed (see abbreviation FT above).
For a numerical answer, allow the A or B mark if the answer is correct to 3 significant figures or would be correct to 3 significant figures if rounded (1 decimal place for angles in degrees).
The total number of marks available for each question is shown at the bottom of the Marks column.
Wrong or missing units in an answer should not result in loss of marks unless the guidance indicates otherwise.
Square brackets [ ] around text or numbers show extra information not needed for the mark to be awarded.

PUBLISHED

Abbreviations

AEF/OE   Any Equivalent Form (of answer is equally acceptable) / Or Equivalent

AG   Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)

CAO   Correct Answer Only (emphasising that no 'follow through' from a previous error is allowed)

CWO   Correct Working Only

ISW   Ignore Subsequent Working

SOI   Seen Or Implied

SC   Special Case (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

WWW   Without Wrong Working

AWRT   Answer Which Rounds To

Question	Answer	Marks	Guidance
1(a)	$\lambda = \dfrac{23.4}{60}$ or 0.39	B1	First **B1** only scored if see 0.39.
	$1 - e^{-0.39}(1 + 0.39) = 1 - (0.67706 + 0.26405)$	M1	Any λ. Allow one end error.
	$= 0.0589$ (3 sf)	A1	**SC** unsupported answer score **B1** instead of **M1A1**.
		3	
1(b)(i)	N(23.4, 23.4)	B1B1	**B1** for N(23.4, ….) **B1** for Var(X) = 23.4. Note: marks for **(i)** cannot be recovered from **(ii)**.
		2	
1(b)(ii)	$\dfrac{20.5 - 23.4}{\sqrt{23.4}}$ [= –0.5995] $\dfrac{29.5 - 23.4}{\sqrt{23.4}}$ [= 1.2610]	M1	Attempt to standardise both with their values. Must have square roots. Allow no (or incorrect) continuity correction(s).
	$\Phi(\text{'}1.2610\text{'}) - \Phi(\text{'}-0.5995\text{'})$ (0.8964 – 0.2743)	M1	For attempt at area between 20 and 30 consistent with their working.
	$= 0.622$ (3 sf)	A1	**SC** no working seen can score **B2** for 0.622.
		3	

Question	Answer	Marks	Guidance
2(a)(i)	$\dfrac{10.03 - 10}{\dfrac{\sigma}{\sqrt{50}}} = 1.995$	M1	For standardising. Must have square root 50.
		M1	Equating to 1.995 (not phi 1.995).
	$= 0.106$ (3 sf)	A1	
		3	

Question	Answer	Marks	Guidance
2(a)(ii)	H$_0$: $\mu = 10$ H$_1$: $\mu > 10$	B1	Accept if seen in part **(i)** but not in **(ii)**. Accept population mean but not just mean.
	1.995 > 1.96	M1	Valid comparison. 0.977 > 0.975 or 0.023 < 0.025. Accept 2.00 or 2.001 > 1.96.
	[Reject H$_0$] There is sufficient evidence [at 2.5% level] to suggest that mean length is greater than 10 cm. Or There is sufft evidence to support the inspectors thought /claim.	A1	No contradictions. In context. Not definite. Accept '….. that mean length has increased' Note 2 tail test scores **B0M1A0**.
		3	
2(b)	No, because population distribution [of lengths] is normal	B1	
		1	

Question	Answer	Marks	Guidance
3(a)(i)	$\text{Est}(\mu) = \dfrac{46350}{150}$ or 309	B1	
	$\text{Est}\left(\sigma^2\right) = \dfrac{150}{149}\left[\dfrac{14410800}{150} - \left(\dfrac{46350}{150}\right)^2\right]\left[=\dfrac{150}{149}\times 591\right]$	M1	Use of correct formula. $\dfrac{1}{149}$ ($14410800 - \dfrac{46350^2}{150}$). OE.
	= 595 (3 sf) or 88650/149	A1	
		3	

Question	Answer	Marks	Guidance
3(a)(ii)	$z = 1.96$	B1	Seen.
	$309 \pm z \times \sqrt{\dfrac{'595'}{150}}$	M1	Any z. Use of correct formula.
	= 305 to 313 (3 sf)	A1	Use of biased (591) gives same interval but scores **B1M1A0**. Unsupported correct answer scores max **B1B1**.
		3	
3(b)	<u>Either</u> Smaller sample means more uncertainty about the value of μ.		OE.
	<u>or</u> $n < 120$ means that $\dfrac{1}{\sqrt{n}} > \dfrac{1}{\sqrt{150}}$	B1*	OE.
	Therefore Wider	B1dep	Dep on previous **B1**.
		2	

Question	Answer	Marks	Guidance
4(a)	The probabilities are not equal	B1	OE. SOI.
	e.g. $P(n=0)=\dfrac{1}{4}$ and $P(n=1)=\dfrac{1}{2}$ or $P(n=1)=2\times P(n=0)$ or similar	B1	Numerical justification or there is one way to get 0 but two ways to get 1, or similar **B1B1**.
		2	

Question	Answer	Marks	Guidance
4(b)	Choose PQ or QR or RP soi or reject R or P or Q soi	B1	Identifying the three correct cases to choose or reject and correctly identifying choosing or rejecting.
	eg 1 or 2 [choose]PQ. 3 or 4 [choose] QR. 5 or 6 [choose]RP		OE.
	or 1 or 2 [reject] R. 3 or 4[reject] P. 5 or 6 [reject] Q	B1	OE. Using the score on the dice to give each of the three cases equal probability. More than one throw of the dice scores **B0B0**.
		2	

Question	Answer	Marks	Guidance
5	Assume the men on the bridge are a random sample	B1	OE. Allow: assume that all the men on the bridge are independent of one another.
	$\dfrac{1500 - 70.3n}{5.9\sqrt{n}} = 2.326$	M1	Must have square root n. Allow wrong z but must be a z value.
	$70.3n + 13.7234\sqrt{n} - 1500 = 0$ Or $4942.09\,n^2 - 211088.33\,n + 2250000 = 0$	A1	Correct quadratic equation or inequality in square root n. OE.
	$[\sqrt{n} = 4.5226$ (3 sf)] Max value of n is 20	A1	CWO.
		4	

Question	Answer	Marks	Guidance
6(a)	$\dfrac{1}{2} \times b \times ab = 1$ or $a\displaystyle\int_0^b x\,dx = 1$	**M1**	Attempt triangle area or integrate f(x) (ignore limits) and = 1.
	$a\left[\dfrac{x^2}{2}\right]_0^b = 1$	**A1**	Correct integration. ½ × b × ab = 1.
	$\dfrac{ab^2}{2} = 1 \Rightarrow a = \dfrac{2}{b^2}$	**A1**	Convincingly obtained. No errors seen.
		3	

Question	Answer	Marks	Guidance
6(b)	$E(X) = \dfrac{2}{b^2} \displaystyle\int_0^b x^2 dx$	M1	Attempt to integrate $xf(x)$. Ignore limits.
	$= \dfrac{2}{b^2} \left[\dfrac{x^3}{3} \right]_0^b$	A1	Correct integration and limits.
	$= \dfrac{2}{3}b$	A1	OE. E.g. $ab^3/3$
	$P(X < E(X)) = \dfrac{2}{b^2} \displaystyle\int_0^{\frac{2}{3}b} x\, dx$	M1	OE. Integrate $f(x)$ with limits 0 and their E(X) Condone a instead of $\dfrac{2}{b^2}$ Or using areas, prob $= \dfrac{1}{2} \times \dfrac{2b}{3} \times \dfrac{2}{3} ab$
	$= \dfrac{2}{b^2} \left[\dfrac{x^2}{2} \right]_0^{\frac{2}{3}b}$	A1FT	Correct integration and limits, **FT** their $E(X)$. Need $\dfrac{2}{b^2}$. Or prob $= \dfrac{1}{2} \times '\dfrac{2b}{3}' \times \dfrac{2}{3} \times \dfrac{2b}{b^2}$.
	$= \dfrac{2}{b^2} \times \dfrac{\frac{4}{9}b^2}{2} = \dfrac{4}{9}$	A1	Must see correct expression after substituting limits. Convincingly obtained. No errors seen. Or expression correctly simplified.
		6	

Question	Answer	Marks	Guidance
7(a)	B(35, 0.25)	**M1**	SOI.
	$0.75^{35} + 35 \times 0.75^{34} \times 0.25 + {}^{35}C_2 \times 0.75^{33} \times 0.25^2 + {}^{35}C_3 \times 0.75^{32} \times 0.25^3$ = 0.000042378 + 0.0004944 + 0.00280168 + 0.01027283	**M1**	Attempt $P(r \leqslant 3)$ Expressions or terms.
	= 0.0136 (3 sf) [< 0.04]	**A1**	**SC** unsupported answer 0.0136 **B1**.
	'0.0136' + ${}^{35}C_4 \times 0.75^{31} \times 0.25^4$ = 0.0136 + 0.0273942	**M1**	Attempt $P(r \leqslant 4)$ Expressions or terms
	= 0.041[0] (2 sf) [> 0.04]	**A1**	**SC** unsupported answer 0.041 **B1**.
	Maximum value of r is 3	**A1**	Dep on above two correct probabilities seen and at least one comparison with 0.04.
		6	
7(b)	Concluding proportion has decreased when in fact it hasn't	**B1**	Or concluding that the director's belief is correct when in fact it isn't.
	0.0136	**B1**	**FT** their $P(r \leqslant 3)$, dep < 0.04 OE.
		2	
7(c)	$1 - (0.95^{35} + 35 \times 0.95^{34} \times 0.05 + {}^{35}C2 \times 0.95^{33} \times 0.05^2 + {}^{35}C3 \times 0.95^{32} \times 0.05^3)$ = 1 − (0.16608 + 0.305943 + 0.2737385 + 0.158480)	**M1**	Attempt $1 - P(r \leqslant 3)$ with B(35, 0.05). Expressions or terms.
	= 0.0958 (3 sf)	**A1**	**SC** unsupported answer 0.0958 **B1**.
		2	