

9.




Figure 3

In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.

Figure 3 shows a sketch of part of the curve C with equation

$$y = \frac{2}{3}x^2 - 9\sqrt{x} + 13 \quad x \geq 0$$

(a) Find, using calculus, the range of values of x for which y is increasing.

(4)

The point P lies on C and has coordinates $(9, 40)$.

The line l is the tangent to C at the point P .

The finite region R , shown shaded in Figure 3, is bounded by the curve C , the line l , the x -axis and the y -axis.

(b) Find, using calculus, the exact area of R .

(8)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT

8.

In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.




Figure 1

Figure 1 shows a sketch of part of the curve C_1 with equation

$$y = x^2 + 3 \quad x > 0$$

and part of the curve C_2 with equation

$$y = 13 - \frac{9}{x^2} \quad x > 0$$

The curves C_1 and C_2 intersect at the points P and Q as shown in Figure 1.(a) Use algebra to find the x coordinate of P and the x coordinate of Q .

(4)

The finite region R , shown shaded in Figure 1, is bounded by C_1 and C_2 (b) Use algebraic integration to find the exact area of R .

(4)

10. The circle C has centre $X(3, 5)$ and radius r

The line l has equation $y = 2x + k$, where k is a constant.

(a) Show that l and C intersect when

$$5x^2 + (4k - 26)x + k^2 - 10k + 34 - r^2 = 0$$

(3)

Given that l is a tangent to C ,

(b) show that $5r^2 = (k + p)^2$, where p is a constant to be found.

(3)




Figure 2

The line l

- cuts the y -axis at the point A
- touches the circle C at the point B

as shown in Figure 2.

Given that $AB = 2r$

(c) find the value of k

(6)

8. In this question you must show all stages of your working.
Solutions relying entirely on calculator technology are not acceptable.




Figure 2

Figure 2 shows a sketch of part of the curve C with equation

$$y = \frac{4}{3}x^3 - 11x^2 + kx \quad \text{where } k \text{ is a constant}$$

The point M is the maximum turning point of C and is shown in Figure 2.

Given that the x coordinate of M is 2

(a) show that $k = 28$ (3)

(b) Determine the range of values of x for which y is increasing. (2)

The line l passes through M and is parallel to the x -axis.

The region R , shown shaded in Figure 2, is bounded by the curve C , the line l and the y -axis.

(c) Find, by algebraic integration, the exact area of R . (5)

8.

In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.




Figure 1

Figure 1 shows a sketch of part of the curve C_1 with equation

$$y = x^2 + 3 \quad x > 0$$

and part of the curve C_2 with equation

$$y = 13 - \frac{9}{x^2} \quad x > 0$$

The curves C_1 and C_2 intersect at the points P and Q as shown in Figure 1.

(a) Use algebra to find the x coordinate of P and the x coordinate of Q .

(4)

The finite region R , shown shaded in Figure 1, is bounded by C_1 and C_2

(b) Use algebraic integration to find the exact area of R .

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT

9.

In this question you must show detailed reasoning.

Solutions relying entirely on calculator technology are not acceptable.




Figure 3

Figure 3 shows

- the curve C_1 with equation $y = x^3 - 5x^2 + 3x + 14$
- the circle C_2 with centre T

The point T is the minimum turning point of C_1

Using Figure 3 and calculus,

(a) find the coordinates of T

(3)

The curve C_1 intersects the circle C_2 at the point A with x coordinate 2

(b) Find an equation of the circle C_2

(3)

The line l shown in Figure 3, is the tangent to circle C_2 at A

(c) Show that an equation of l is

$$y = \frac{1}{3}x + \frac{22}{3}$$

(3)

The region R , shown shaded in Figure 3, is bounded by C_1 , l and the y -axis.

(d) Find the exact area of R .

(3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

6. A circle has equation

$$x^2 - 6x + y^2 + 8y + k = 0$$

where k is a positive constant.

Given that the x -axis is a tangent to this circle,

(a) find the value of k .

(3)

The circle meets the coordinate axes at the points R , S and T .

(b) Find the exact area of the triangle RST .

(4)

9.




Figure 2

Figure 2 shows

- the curve C with equation $y = x - x^2$
- the line l with equation $y = mx$, where m is a constant and $0 < m < 1$

The line and the curve intersect at the origin O and at the point P .

(a) Find, in terms of m , the coordinates of P .

(2)

The region R_1 , shown shaded in Figure 2, is bounded by C and l .

(b) Show that the area of R_1 is

$$\frac{(1-m)^3}{6}$$

(5)

The region R_2 , also shown shaded in Figure 2, is bounded by C , the x -axis and l .

Given that the area of R_1 is equal to the area of R_2

(c) find the exact value of m .

(3)


blank

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

6.

**Figure 2**

The points $A(-2, 18)$ and $B(14, 6)$ lie on a circle C_1 as shown in Figure 2.

Given that AB is a diameter of the circle C_1

(a) find an equation for C_1 making your method clear.

(5)

A circle C_2 has its centre at the origin.

Given that circles C_1 and C_2 touch,

(b) find possible equations for C_2

(4)

7.




Figure 3

In this question you must show all stages of your working.
Solutions relying entirely on calculator technology are not acceptable.

Figure 3 shows a sketch of part of the curve C with equation

$$y = x^3 - 4x^{\frac{5}{2}} - kx^{\frac{1}{2}} + 28x - 44 \quad x \geq 0$$

where k is a positive constant.

(a) Find $\frac{dy}{dx}$ in simplest form.

(2)

The point T , shown in Figure 3, is a minimum stationary point on C .

Given that the x coordinate of T is 9

(b) show that $k = 6$

(2)

The line through T parallel to the x -axis meets the y -axis at the point N .

The finite region R , shown shaded in Figure 3, is bounded by C , the y -axis and the line segment NT .

(c) Use algebraic integration to find the area of R , giving the answer to 3 significant figures.

(6)

9. A circle C has equation

$$(x - k)^2 + (y - 2k)^2 = k + 7$$

where k is a positive constant.

(a) Write down, in terms of k ,

(i) the coordinates of the centre of C ,

(ii) the radius of C .

(2)

Given that the point $P(2, 3)$ lies on C

(b) (i) show that $5k^2 - 17k + 6 = 0$

(ii) hence find the possible values of k .

(3)

The tangent to the circle at P intersects the x -axis at point T .

Given that $k < 2$

(c) calculate the exact area of triangle OPT .

(5)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

6 It is given that $3 \sin 2\theta = \cos \theta$ where θ is an angle such that $0^\circ < \theta < 90^\circ$.

(a) Find the exact value of $\sin \theta$.

[2]

.....
.....
.....
.....
.....
.....

(b) Find the exact value of $\sec \theta$.

[2]

.....
.....
.....
.....
.....
.....

(c) Find the exact value of $\cos 2\theta$.

[2]

10:




Figure 3

**In this question you must show all stages of your working.
Solutions relying entirely on calculator technology are not acceptable.**

Figure 3 shows a sketch of the curve C with equation

$$y = 2x + \frac{64}{x^2} - 3 \quad x > 0$$

The point P , shown in Figure 3, is the stationary point on C .

(a) Show, using calculus, that the x coordinate of P is 4

(4)

The point Q lies on C and has x coordinate 2

The line segments MQ and NP , shown in Figure 3, are parallel to the x -axis.

The region R , shown shaded in Figure 3, is bounded by C , the y -axis and line segments MQ and NP .

(b) Use algebraic integration to find the exact area of R .

You must make your method clear.

(5)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA