4. The function f is defined by

$$f(x) = 2x^2 - 5 \qquad x \geqslant 0 \quad x \in \mathbb{R}$$

(a) State the range of f

(1)

On the following page there is a diagram, labelled Diagram 1, which shows a sketch of the curve with equation $y = f(x)$.

(b) On Diagram 1, sketch the curve with equation $y = f^{-1}(x)$.

(2)

The curve with equation $y = f(x)$ meets the curve with equation $y = f^{-1}(x)$ at the point P

Using algebra and showing your working,

(c) find the exact x coordinate of P

(3)

3. (i) The variables x and y are connected by the equation

$$y = \frac{10^6}{x^3} \qquad x > 0$$

Sketch the graph of $\log_{10} y$ against $\log_{10} x$

Show on your sketch the coordinates of the points of intersection of the graph with the axes.

(3)

(ii)



Figure 2

Figure 2 shows the linear relationship between $\log_3 N$ and t.

Show that $N = ab^t$ where a and b are constants to be found.

(3)

7. **In this question you must show all stages of your working.**

Solutions relying entirely on calculator technology are not acceptable.

The curve C has equation

$$y = \frac{16}{9(3x - k)} \qquad x \neq \frac{k}{3}$$

where k is a positive constant not equal to 3

(a) Find $\frac{dy}{dx}$ giving your answer in simplest form in terms of k.

(2)

The point P with x coordinate 1 lies on C.

Given that the gradient of the curve at P is -12

(b) find the two possible values of k

(3)

Given also that $k < 3$

(c) find the equation of the normal to C at P, writing your answer in the form $ax + by + c = 0$, where a, b and c are integers to be found.

(3)

(d) show, using algebraic integration that,

$$\int_1^3 \frac{16}{9(3x - k)} dx = \lambda \ln 10$$

where λ is a constant to be found.

(4)

4. $$\mathrm{f}(x) = 8\sin x\cos x + 4\cos^2 x - 3$$

(a) Write $\mathrm{f}(x)$ in the form

$$a\sin 2x + b\cos 2x + c$$

where a, b and c are integers to be found.

(3)

(b) Use the answer to part (a) to write $\mathrm{f}(x)$ in the form

$$R\sin(2x + \alpha) + c$$

where $R > 0$ and $0 < \alpha < \frac{\pi}{2}$

Give the exact value of R and give the value of α in radians to 3 significant figures.

(3)

(c) Hence, or otherwise,

(i) state the maximum value of $\mathrm{f}(x)$

(ii) find the **second** smallest positive value of x at which a maximum value of $\mathrm{f}(x)$ occurs. Give your answer to 3 significant figures.

(3)