

7.




Figure 3

The curve C has equation $y = f(x)$, where

$$f(x) = x^3 \sqrt{4x + 7} \quad x \geq -\frac{7}{4}$$

(a) Show that

$$f'(x) = \frac{kx^2(2x + 3)}{\sqrt{4x + 7}}$$

where k is a constant to be found.

(4)

The point P , shown in Figure 3, is the minimum turning point on C .

(b) Find the coordinates of P .

(2)

(c) Hence find the range of the function g defined by

$$g(x) = -4f(x) \quad x \geq -\frac{7}{4}$$

(2)

The point Q with coordinates $\left(\frac{1}{2}, \frac{3}{8}\right)$ lies on C .

(d) Find the coordinates of the point to which Q is mapped when C is transformed to the curve with equation

$$y = 40f\left(x - \frac{3}{2}\right) - 8$$

(2)

9.




Figure 5

The curve shown in Figure 5 has equation

$$x = 4\sin^2 y - 1 \quad 0 \leq y \leq \frac{\pi}{2}$$

The point $P\left(k, \frac{\pi}{3}\right)$ lies on the curve.

(a) Verify that $k = 2$ (1)

(b) (i) Find $\frac{dx}{dy}$ in terms of y

(ii) Hence show that $\frac{dy}{dx} = \frac{1}{2\sqrt{x+1}\sqrt{3-x}}$ (6)

The normal to the curve at P cuts the x -axis at the point N .

(c) Find the exact area of triangle OPN , where O is the origin.

Give your answer in the form $a\pi + b\pi^2$ where a and b are constants. (3)

7.

In this question you must show all stages of your working.**Solutions relying entirely on calculator technology are not acceptable.**The curve C has equation

$$y = \frac{16}{9(3x - k)} \quad x \neq \frac{k}{3}$$

where k is a positive constant not equal to 3

- (a) Find $\frac{dy}{dx}$ giving your answer in simplest form in terms of k .

(2)

The point P with x coordinate 1 lies on C .Given that the gradient of the curve at P is -12

- (b) find the two possible values of k

(3)

Given also that $k < 3$

- (c) find the equation of the normal to C at P , writing your answer in the form $ax + by + c = 0$, where a , b and c are integers to be found.

(3)

- (d) show, using algebraic integration that,

$$\int_1^3 \frac{16}{9(3x - k)} dx = \lambda \ln 10$$

where λ is a constant to be found.

(4)

10.




Figure 4

Figure 4 shows a sketch of the curve with equation

$$x = \frac{2y^2 + 6}{3y - 3}$$

- (a) Find $\frac{dx}{dy}$ giving your answer as a fully simplified fraction.

(4)

The tangents at points P and Q on the curve are parallel to the y -axis, as shown in Figure 4.

- (b) Use the answer to part (a) to find the equations of these two tangents.

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

2.




Figure 1

Figure 1 shows a sketch of the curve with equation

$$x = 2y^2 + 5y - 6$$

- (a) Find $\frac{dy}{dx}$ in terms of y .

(2)

The point P lies on the curve and is shown in Figure 1.

Given that the tangent to the curve at P is parallel to the y -axis,

- (b) find the coordinates of P .

(3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

8.




Figure 3

Figure 3 shows a sketch of the curve C with equation $y = f(x)$, where

$$f(x) = (2x + 1)^3 e^{-4x}$$

(a) Show that

$$f'(x) = A(2x + 1)^2 (1 - 4x) e^{-4x}$$

where A is a constant to be found.

(4)

(b) Hence find the exact coordinates of the two stationary points on C .

(3)

The function g is defined by

$$g(x) = 8f(x - 2)$$

(c) Find the coordinates of the maximum stationary point on the curve with equation $y = g(x)$.

(2)

10.

In this question you must show all stages of your working.**Solutions relying on calculator technology are not acceptable.**A curve C has equation

$$x = \sin^2 4y \quad 0 \leq y \leq \frac{\pi}{8} \quad 0 \leq x \leq 1$$

The point P with x coordinate $\frac{1}{4}$ lies on C (a) Find the exact y coordinate of P

(2)

(b) Find $\frac{dx}{dy}$

(2)

(c) Hence show that $\frac{dy}{dx}$ can be written in the form

$$\frac{dy}{dx} = \frac{1}{\sqrt{q + r(x + s)^2}}$$

where q , r and s are constants to be found.

(3)

Using the answer to part (c),

(d) (i) state the x coordinate of the point where the value of $\frac{dy}{dx}$ is a minimum,(ii) state the value of $\frac{dy}{dx}$ at this point.

(2)

5:

In this question you must show all stages of your working.
Solutions relying entirely on calculator technology are not acceptable.

A curve C has equation

$$y = 5 \cos 2x - 12 \sin 2x \quad -\pi < x < \pi$$

The point A with x coordinate $\frac{\pi}{3}$ lies on C .

- (a) Use algebraic differentiation to find the gradient of the tangent to C at A .
Give the answer in simplest form.

(3)

- (b) Express $5 \cos 2x - 12 \sin 2x$ in the form

$$R \cos(2x + \alpha)$$

where $R > 0$ and $0 < \alpha < \frac{\pi}{2}$

Find the exact value of R and find the value of α to 3 decimal places.

(3)

A point P lies on C .

Given that the gradient of the tangent to C at P is 6

- (c) find the greatest possible value for the x coordinate of P .
Give the answer to 2 decimal places.

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

6.




Figure 3

In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.

Figure 3 shows a sketch of part of the curve with equation

$$y = \sqrt{4x - 7}$$

The line l , shown in Figure 3, is the normal to the curve at the point $P(8, 5)$

(a) Use calculus to show that an equation of l is

$$5x + 2y - 50 = 0 \quad (5)$$

The region R , shown shaded in Figure 3, is bounded by the curve, the x -axis and l .

(b) Use algebraic integration to find the exact area of R .

(4)

9.


**Figure 1**

Figure 1 shows a sketch of part of the curve C with equation $y = f(x)$ where

$$f(x) = 6\sqrt{x} \ln(4x) \quad x > 0$$

The curve cuts the x -axis at point P

- (a) State the x coordinate of P

(1)

The point Q , shown in Figure 1, is the stationary point on C

- (b) Use calculus to find the exact coordinates of Q

(5)

- (c) Hence find the range of the function $g(x)$ where

$$g(x) = -2f(x)$$

(2)

3.

In this question you must show all stages of your working.

Solutions relying on calculator technology are not acceptable.

A curve has equation

$$y = \frac{4x+1}{(x+3)^2} \quad x \neq -3 \quad x \in \mathbb{R}$$

Use calculus to find the range of values of x for which y is increasing.

(6)

4: The function f is defined by

$$f(x) = \frac{49x}{x^2 + x - 12} + \frac{7x}{x + 4} \quad x > 3$$

(a) Show that

$$f(x) = \frac{7x}{x - 3} \quad x > 3 \quad (3)$$

(b) Hence find $f'(x)$ giving your answer in simplest form.

(2)

5. The curve C has equation

$$y = \frac{\ln(x^2 + k)}{x^2 + k} \quad x \in \mathbb{R}$$

where k is a positive constant.

(a) Show that

$$\frac{dy}{dx} = \frac{Ax(B - \ln(x^2 + k))}{(x^2 + k)^2} \quad (3)$$

where A and B are constants to be found.

Given that C has exactly three turning points,

(b) find the x coordinate of each of these points. Give your answer in terms of k where appropriate.

(3)

(c) find the upper limit to the value for k .

(1)

6.




Figure 1

In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.

Figure 1 shows a sketch of the curve with equation $y = f(x)$, where

$$f(x) = 2e^{3\sin x} \cos x \quad 0 \leq x \leq 2\pi$$

The curve intersects the x -axis at point R , as shown in Figure 1.

(a) State the coordinates of R

(1)

The curve has two turning points, at point P and point Q , also shown in Figure 1.

(b) Show that, at points P and Q ,

$$a \sin^2 x + b \sin x + c = 0$$

where a , b and c are integers to be found.

(4)

(c) Hence find the x coordinate of point Q , giving your answer to 3 decimal places.

(2)

9:




Figure 2

Figure 2 shows a sketch of the curve C with equation

$$x = \frac{2}{3} \sin\left(3y + \frac{\pi}{4}\right) \quad \frac{\pi}{12} < y < \frac{3\pi}{4}$$

The curve intersects the y -axis at the points A and B as shown.

(a) Find the exact value of the y coordinate of

- point A
- point B

(3)

(b) Show that

$$\left(\frac{dy}{dx}\right)^2 = \frac{1}{p - qx^2}$$

where p and q are integers to be found.

(4)

The **normal** to C at A and the **tangent** to C at B intersect at the point D .

Using

- the answer to part (b)
- the sketch of curve C in Figure 2

(c) find, in simplest form, the exact x coordinate of D .

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

6:




Figure 2

Figure 2 shows a sketch of the curve with equation $y = f(x)$ where

$$f(x) = (2x - 3)^3 e^{4x-2}$$

(a) Show that

$$f'(x) = 2(Px + Q)(2x - 3)^n e^{4x-2}$$

where P , Q and n are constants to be found.

(4)

(b) Hence find

- (i) the x coordinates of the stationary points on the curve with equation $y = f(x)$,
- (ii) the range of the function g defined by

$$g(x) = 8(2x - 3)^3 e^{4x-2} \quad x \leq \frac{3}{2}$$

(Solutions relying entirely on calculator technology are not acceptable.)

(4)

7.




Figure 1

Figure 1 shows a sketch of the curve C with equation $y = f(x)$ where

$$f(x) = e^{-x^2} (2x^2 - 3)^2$$

(a) Find the range of f

(2)

(b) Show that

$$f'(x) = 2x(2x^2 - 3)e^{-x^2}(A - Bx^2)$$

where A and B are constants to be found.

(4)

Given that the line $y = k$, where k is a constant, $k > 0$, intersects the curve at exactly two distinct points,

(c) find the exact range of values of k

(4)

7.

**In this question you must show all stages of your working.
Solutions relying entirely on calculator technology are not acceptable.**

A curve C has equation

$$y = 2xe^{x^2 + (3k-2)x}$$

where k is a constant.

Given that C has two distinct turning points, find the range of possible values of k .

(7)

9.




Figure 3

Figure 3 shows a sketch of part of the curve C with equation

$$y = \sqrt{3 + 4e^{x^2}} \quad x \geq 0$$

- (a) Find $\frac{dy}{dx}$, giving your answer in simplest form.

(2)

The point P with x coordinate α lies on C .

Given that the tangent to C at P passes through the origin, as shown in Figure 3,

- (b) show that $x = \alpha$ is a solution of the equation

$$4x^2e^{x^2} - 4e^{x^2} - 3 = 0$$

(3)

- (c) Hence show that α lies between 1 and 2

(2)

- (d) Show that the equation in part (b) can be written in the form

$$x = \frac{1}{2}\sqrt{4 + 3e^{-x^2}}$$

(1)

The iteration formula

$$x_{n+1} = \frac{1}{2}\sqrt{4 + 3e^{-x_n^2}}$$

with $x_1 = 1$ is used to find an approximation for α .

- (e) Use the iteration formula to find, to 4 decimal places, the value of

(i) x_3

(ii) α

6.

In this question you must show all stages of your working.
Solutions relying entirely on calculator technology are not acceptable.

(i) Given $y = \ln(2x^2 + 5)$ find $\frac{dy}{dx}$

(2)

(ii) A curve C has equation $y = f(x)$ where

$$f(x) = \frac{21x}{3x^2 + k} \quad x \in \mathbb{R}$$

where k is a positive integer, $k > 1$

Given that

$$\int_1^k f(x) \, dx < 7 \ln 8$$

find the greatest possible value of k

(5)

10.




Figure 2

Figure 2 shows a sketch of the curve with equation

$$x = 3 \cos 2y \quad -3 \leq x \leq 3 \quad 0 \leq y \leq \frac{\pi}{2}$$

(a) Find $\frac{dx}{dy}$ in terms of y .

(2)

(b) Hence show that

$$\frac{dy}{dx} = \frac{k}{\sqrt{9 - x^2}}$$

where k is a constant to be found.

(3)

The point $P(a, b)$ lies on the curve and is shown in Figure 2.

Given that

- the gradient of the curve at P is $-\frac{1}{4}$
- both a and b are positive

(c) find the exact values of a and b .

(4)

3: The curve C has equation

$$x = \frac{4y^2 - 3}{2y + 1} \quad y \neq -\frac{1}{2}$$

(a) Find $\frac{dy}{dx}$ in terms of y , giving the answer in simplest form.

(4)

A point P lies on C .

Given that

- the gradient of the tangent to C at P is $\frac{1}{3}$
- the point P lies above the x -axis

(b) find the coordinates of P .

(Solutions relying entirely on calculator technology are not acceptable.)

(4)

9.

In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.

(a) Show that

$$\frac{\cos 2x}{\sin x} + \frac{\sin 2x}{\cos x} \equiv \operatorname{cosec} x \quad x \neq \frac{n\pi}{2} \quad n \in \mathbb{Z}$$

(3)

(b) Hence solve, for $0 < \theta < \frac{\pi}{2}$

$$\left(\frac{\cos 2\theta}{\sin \theta} + \frac{\sin 2\theta}{\cos \theta} \right)^2 = 6 \cot \theta - 4$$

giving your answers to 3 significant figures as appropriate.

(5)

(c) Using the result from part (a), or otherwise, find the exact value of

$$\int_{\frac{\pi}{6}}^{\frac{\pi}{4}} \left(\frac{\cos 2x}{\sin x} + \frac{\sin 2x}{\cos x} \right) \cot x \, dx$$

(2)