10.



Figure 4

Figure 4 shows a sketch of the curve with equation

$$x = \frac{2y^2 + 6}{3y - 3}$$

(a) Find $\frac{dx}{dy}$ giving your answer as a fully simplified fraction.

(4)

The tangents at points P and Q on the curve are parallel to the y-axis, as shown in Figure 4.

(b) Use the answer to part (a) to find the equations of these two tangents.

(4)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

5. The curve C has equation

$$y = \frac{\ln(x^2 + k)}{x^2 + k} \qquad x \in \mathbb{R}$$

where k is a positive constant.

(a) Show that

$$\frac{dy}{dx} = \frac{Ax(B - \ln(x^2 + k))}{(x^2 + k)^2}$$

where A and B are constants to be found.

(3)

Given that C has exactly three turning points,

(b) find the x coordinate of each of these points. Give your answer in terms of k where appropriate.

(3)

(c) find the upper limit to the value for k.

(1)

---

7. **In this question you must show all stages of your working.**
Solutions relying entirely on calculator technology are not acceptable.

A curve C has equation

$$y = 2xe^{x^2 + (3k-2)x}$$

where k is a constant.

Given that C has two distinct turning points, find the range of possible values of k.

(7)

DO NOT WRITE IN THIS AREA DO NOT WRITE IN THIS

2.



Figure 1

Figure 1 shows a sketch of the curve with equation

$$x = 2y^2 + 5y - 6$$

(a) Find $\frac{dy}{dx}$ in terms of y.

(2)

The point P lies on the curve and is shown in Figure 1.

Given that the tangent to the curve at P is parallel to the y-axis,

(b) find the coordinates of P.

(3)

4.

$$f(x) = 8\sin x \cos x + 4\cos^2 x - 3$$

(a) Write $f(x)$ in the form

$$a\sin 2x + b\cos 2x + c$$

where a, b and c are integers to be found.

(3)

(b) Use the answer to part (a) to write $f(x)$ in the form

$$R\sin(2x + \alpha) + c$$

where $R > 0$ and $0 < \alpha < \frac{\pi}{2}$

Give the exact value of R and give the value of α in radians to 3 significant figures.

(3)

(c) Hence, or otherwise,

(i) state the maximum value of $f(x)$

(ii) find the **second** smallest positive value of x at which a maximum value of $f(x)$ occurs. Give your answer to 3 significant figures.

(3)

DO NOT WRITE IN THIS AREA DO NOT WRITE IN THIS AREA DO NOT WRITE IN THIS AREA DO NOT WR

2: **In this question you must show all stages of your working.**
Solutions relying entirely on calculator technology are not acceptable.

$$f(x) = 7\cos x - 24\sin x$$

(a) Express $f(x)$ in the form $R\cos(x + \alpha)$ where R and α are constants, $R > 0$

and $0 < \alpha < \frac{\pi}{2}$

Give the exact value of R and give the value of α, in radians, to 3 decimal places.

(3)

$$g(x) = \frac{5}{90 - 3f(2x)}$$

(b) Using the answer to part (a), find

(i) the minimum value of $g(x)$, giving your answer as a fully simplified fraction,

(ii) the smallest positive value of x for which this minimum value occurs, giving your answer to 3 decimal places.

(4)

DO NOT WRITE IN THIS AREA DO NOT

7. **In this question you must show all stages of your working.**

Solutions relying entirely on calculator technology are not acceptable.

The curve C has equation

$$y = \frac{16}{9(3x - k)} \qquad x \neq \frac{k}{3}$$

where k is a positive constant not equal to 3

(a) Find $\frac{dy}{dx}$ giving your answer in simplest form in terms of k.

(2)

The point P with x coordinate 1 lies on C.

Given that the gradient of the curve at P is -12

(b) find the two possible values of k

(3)

Given also that $k < 3$

(c) find the equation of the normal to C at P, writing your answer in the form $ax + by + c = 0$, where a, b and c are integers to be found.

(3)

(d) show, using algebraic integration that,

$$\int_1^3 \frac{16}{9(3x - k)} dx = \lambda \ln 10$$

where λ is a constant to be found.

(4)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

5: **In this question you must show all stages of your working.**
Solutions relying entirely on calculator technology are not acceptable.

A curve C has equation

$$y = 5\cos 2x - 12\sin 2x \qquad -\pi < x < \pi$$

The point A with x coordinate $\frac{\pi}{3}$ lies on C.

(a) Use algebraic differentiation to find the gradient of the tangent to C at A.
Give the answer in simplest form.

(3)

(b) Express $5\cos 2x - 12\sin 2x$ in the form

$$R\cos(2x + \alpha)$$

where $R > 0$ and $0 < \alpha < \frac{\pi}{2}$

Find the exact value of R and find the value of α to 3 decimal places.

(3)

A point P lies on C.

Given that the gradient of the tangent to C at P is 6

(c) find the greatest possible value for the x coordinate of P.
Give the answer to 2 decimal places.

(4)

6. **In this question you must show all stages of your working.**
Solutions relying entirely on calculator technology are not acceptable.

(i) Given $y = \ln(2x^2 + 5)$ find $\frac{dy}{dx}$

(2)

(ii) A curve C has equation $y = f(x)$ where

$$f(x) = \frac{21x}{3x^2 + k} \qquad x \in \mathbb{R}$$

where k is a positive integer, $k > 1$

Given that

$$\int_1^k f(x)\,dx < 7\ln 8$$

find the greatest possible value of k

(5)

DO NOT WRITE IN THIS AREA DO NOT WRITE IN THIS AREA DO NOT WRITE IN THIS AREA

1. **In this question you must show all stages of your working.**

Solutions relying entirely on calculator technology are not acceptable.

Solve, for $0 < \theta \leqslant 360°$, the equation

$$3\tan^2\theta + 7\sec\theta - 3 = 0$$

giving your answers to one decimal place.

(5)

DO NOT WRITE IN THIS