

9.




Figure 4

Figure 4 shows a sketch of the curve C with parametric equations

$$x = \sec t \quad y = \sqrt{3} \tan\left(t + \frac{\pi}{3}\right) \quad \frac{\pi}{6} < t < \frac{\pi}{2}$$

(a) Find $\frac{dy}{dx}$ in terms of t

(3)

(b) Find an equation for the tangent to C at the point where $t = \frac{\pi}{3}$

Give your answer in the form $y = mx + c$, where m and c are constants.

(4)

(c) Show that all points on C satisfy the equation

$$y = \frac{Ax^2 + B\sqrt{3x^2 - 3}}{4 - 3x^2}$$

where A and B are constants to be found.

(5)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

EA

DO

8.




Figure 2

Figure 2 shows a sketch of part of the curve C with parametric equations

$$x = t + \frac{1}{t} \quad y = t - \frac{1}{t} \quad t > 0.7$$

The curve C intersects the x -axis at the point Q .

(a) Find the x coordinate of Q .

(1)

The line l is the normal to C at the point P as shown in Figure 2.

Given that $t = 2$ at P

(b) write down the coordinates of P

(1)

(c) Using calculus, show that an equation of l is

$$3x + 5y = 15$$

(3)

The region, R , shown shaded in Figure 2 is bounded by the curve C , the line l and the x -axis.

(d) Using algebraic integration, find the exact volume of the solid of revolution formed when the region R is rotated through 2π radians about the x -axis.

(7)

8.

$$f(x) = (8 - 3x)^{\frac{4}{3}} \quad 0 < x < \frac{8}{3}$$

- (a) Show that the binomial expansion of $f(x)$ in ascending powers of x up to and including the term in x^3 is

$$A - 8x + \frac{x^2}{2} + Bx^3 + \dots$$

where A and B are constants to be found.

(4)

- (b) Use proof by contradiction to prove that the curve with equation

$$y = 8 + 8x - \frac{15}{2}x^2$$

does **not** intersect the curve with equation

$$y = A - 8x + \frac{x^2}{2} + Bx^3 \quad 0 < x < \frac{8}{3}$$

where A and B are the constants found in part (a).

(Solutions relying on calculator technology are not acceptable.)

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

5.




Figure 2

Figure 2 shows a sketch of the curve C with equation

$$y^3 - x^2 + 4x^2y = k$$

where k is a positive constant greater than 1

(a) Find $\frac{dy}{dx}$ in terms of x and y .

(5)

The point P lies on C .

Given that the normal to C at P has equation $y = x$, as shown in Figure 2,

(b) find the value of k .

(5)




Figure 2

In this question you must show all stages of your working.
Solutions relying entirely on calculator technology are not acceptable.

Figure 2 shows a sketch of the curve C with parametric equations

$$x = 2 \cos 2t \quad y = \sin^3 t \quad -\frac{\pi}{2} < t < \frac{\pi}{2}$$

where t is a parameter.

The point P lies on C where $t = \frac{\pi}{6}$

The line l , shown in Figure 2, is the tangent to C at P .

(a) Use parametric differentiation to show that

(i) $\frac{dy}{dx} = k \sin t$ where k is a constant to be found

(ii) an equation for l is $3x + 16y - 5 = 0$

(6)

The line l intersects the curve C again at the point Q .

(b) Using algebra and showing detailed reasoning, find the exact coordinates of Q .

(6)

3: Given that $y = 0$ at $x = 5$ solve the differential equation

$$8y \frac{dy}{dx} = xe^{-y^2} \quad y \geq 0 \quad x \geq 5$$

giving your answer in the form $y^2 = f(x)$

(5)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA